

Asymptotic Expansions for Stationary Distributions of Nonlinearly Perturbed Semi-Markov Processes. II

Dmitrii Silvestrov¹ and Sergei Silvestrov²

Abstract: Asymptotic expansions with explicit upper bounds for remainders are given for stationary distributions of nonlinearly perturbed semi-Markov processes with finite phase spaces. The corresponding algorithms are based on a special technique of sequential phase space reduction, which can be applied to processes with an arbitrary asymptotic communicative structure of phase spaces.

Keywords: Markov chain; semi-Markov process; nonlinear perturbation; stationary distribution; expected hitting time; Laurent asymptotic expansion

2010 Mathematics Subject Classification: Primary 60J10, 60J27, 60K15, Secondary 65C40.

1. Introduction

In this paper, we present new algorithms for construction of asymptotic expansions for stationary distributions of nonlinearly perturbed semi-Markov processes with a finite phase space. This is Part II of the paper, Silvestrov, D. and Silvestrov, S. (2016), where algorithms for constructing of asymptotic expansions with remainders of a standard form $o(\cdot)$ have been given. In Part II, we present algorithms for construction asymptotic expansions of a more advanced form, with explicit upper bounds for remainders.

We consider models, where the phase space is one class of communicative states, for embedded Markov chains of pre-limiting perturbed semi-Markov processes, while it can possess an arbitrary communicative structure, i.e., can

¹Department of Mathematics, Stockholm University, SE-106 81 Stockholm, Sweden.
Email address: silvestrov@math.su.se

²Division of Applied Mathematics, School of Education, Culture and Communication, Mälardalen University, SE-721 23 Västerås, Sweden.
Email address: sergei.silvestrov@mdh.se

consist of one or several closed classes of communicative states and, possibly, a class of transient states, for the limiting embedded Markov chain.

The initial perturbation conditions are formulated in the forms of Taylor and Laurent asymptotic expansions with explicit upper bounds for remainders, respectively, for transition probabilities (of embedded Markov chains) and expectations of sojourn times, for perturbed semi-Markov processes.

The algorithms are based on special time-space screening procedures for sequential phase space reduction and algorithms for re-calculation of asymptotic expansions with explicit upper bounds for remainders, which constitute perturbation conditions for the semi-Markov processes with reduced phase spaces.

The final asymptotic expansions for stationary distributions of nonlinearly perturbed semi-Markov processes are given in the form of Taylor asymptotic expansions, with explicit upper bounds for remainders.

The algorithms presented in the paper have an universal character. They can be applied to perturbed semi-Markov processes with an arbitrary asymptotic communicative structure and are computationally effective due to recurrent character of computational procedures.

The survey of works in the area and detailed comments concerned with the proposed method are given in the Introduction to Part I of this paper.

Here, we would like, only, to mention books, which contains parts devoted to perturbed Markov chains and semi-Markov processes, and problems related to asymptotic expansions for the above models. These are, Korolyuk and Turbin (1976, 1978), Courtois (1977), Seneta (1981, 2006), Stewart and Sun (1990), Kartashov (1996), Stewart (1998, 2001), Yin and Zhang (1998, 2005, 2013), Korolyuk, V.S. and Korolyuk, V.V. (1999), Konstantinov, Gu, Mehrmann and Petkov (2003), Bini, Latouche and Meini (2005), Korolyuk and Limnios (2005), Gyllenberg and Silvestrov (2008), and Avrachenkov, Filar and Howlett (2013).

A comprehensive bibliography of works in the area can be found in these books and, also, in the research report by Silvestrov, D. and Silvestrov, S. (2015), which is an extended preliminary version of the present paper.

In conclusion, we would like to mention that, by our opinion, the results presented in the paper have a good potential for continuation of studies. We comment some prospective directions for future studies in the last section of the paper.

Part II includes four sections and two appendices. In Section 2, we present so-called operational rules for Laurent asymptotic expansions with explicit

upper bounds for remainders. In Section 3, we present basic perturbation conditions and algorithms for construction of asymptotic expansions with explicit upper bounds for remainders, for transition characteristics of nonlinearly perturbed semi-Markov processes with reduced phase spaces. In Section 4, we present algorithms for construction of asymptotic expansions with explicit upper bounds for remainders, for expected hitting times for nonlinearly perturbed semi-Markov processes. In Section 5, we present an algorithm for construction of asymptotic expansions with explicit upper bounds for remainders, for stationary distributions of nonlinearly perturbed semi-Markov processes. In Appendix A, we give proofs of lemmas representing operational rules for Laurent asymptotic expansions without and with explicit upper bounds for remainders. In Appendix B, we discuss and present examples illustrating algorithms for construction of asymptotic expansions for stationary distributions of nonlinearly perturbed semi-Markov processes developed in the present paper.

2. Laurent asymptotic expansions with explicit upper bounds for remainders

In this section, we present so-called operational rules for Laurent asymptotic expansions with explicit upper bounds for remainders. The corresponding proofs and comments are given in Appendix A.

Let $A(\varepsilon)$ be a real-valued function defined on an interval $(0, \varepsilon_0]$, for some $0 < \varepsilon_0 \leq 1$, and given on this interval by a Laurent asymptotic expansion,

$$A(\varepsilon) = a_{h_A} \varepsilon^{h_A} + \dots + a_{k_A} \varepsilon^{k_A} + o_A(\varepsilon^{k_A}), \quad (1)$$

where **(a)** $-\infty < h_A \leq k_A < \infty$ are integers, **(b)** coefficients $a_{h_A}, \dots, a_{k_A}$ are real numbers, **(c)** $|o_A(\varepsilon^{k_A})| \leq G_A \varepsilon^{k_A + \delta_A}$, for $0 < \varepsilon \leq \varepsilon_A$, where **(d)** $0 < \delta_A \leq 1, 0 < G_A < \infty$ and $0 < \varepsilon_A \leq \varepsilon_0$.

We refer to such Laurent asymptotic expansion as a $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion.

The $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion is also a (h_A, k_A) -expansion, according the definition given in Part I of the paper, since, $o_A(\varepsilon^{k_A})/\varepsilon^{k_A} \rightarrow 0$ as $\varepsilon \rightarrow 0$.

We say that $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion $A(\varepsilon)$ is pivotal if it is known that $a_{h_A} \neq 0$.

It is useful to note that there is no sense to consider, it seems, a more general case of upper bounds for the remainder $o_A(\varepsilon^{k_A})$, with parameter $\delta_A >$

1. Indeed, let us define $k'_A = k_A + [\delta_A] - \mathbf{I}(\delta_A = [\delta_A])$ and $\delta'_A = \delta_A - [\delta_A] + \mathbf{I}(\delta_A = [\delta_A]) \in (0, 1]$. The $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion $A(\varepsilon)$ can be re-written in the equivalent form of the $(h_A, k'_A, \delta'_A, G_A, \varepsilon_A)$ -expansion, $A(\varepsilon) = a_{h_A} \varepsilon^{h_A} + \dots + a_{k_A} \varepsilon^{k_A} + 0\varepsilon^{k_A+1} + \dots + 0\varepsilon^{k'_A} + o'_A(\varepsilon^{k'_A})$, with the remainder $o'_A(\varepsilon^{k'_A}) = o_A(\varepsilon^{k_A})$, which satisfies inequalities $|o'_A(\varepsilon^{k'_A})| = |o_A(\varepsilon^{k_A})| \leq G_A \varepsilon^{k_A + \delta_A} = G_A \varepsilon^{k'_A + \delta'_A}$, for $0 < \varepsilon \leq \varepsilon_A$.

The above remarks imply that the asymptotic expansion $A(\varepsilon)$ can be represented in different forms. In such cases, we consider forms with larger parameters h_A and k_A as more informative. As far as parameters δ_A, G_A and ε_A are concerned, we consider as more informative forms, first, with larger values of parameter δ_A , second, with smaller values of parameter G_A and, third, with larger values of parameter ε_A .

In what follows, lemmas, theorems and relations from Part I of the paper are indexed by symbol $*$.

The following proposition supplements Lemma 1 $_*$.

Lemma 1. *If $A(\varepsilon) = a'_{h'_A} \varepsilon^{h'_A} + \dots + a'_{k'_A} \varepsilon^{k'_A} + o'_A(\varepsilon^{k'_A}) = a''_{h''_A} \varepsilon^{h''_A} + \dots + a''_{k''_A} \varepsilon^{k''_A} + o''_A(\varepsilon^{k''_A})$, $\varepsilon \in (0, \varepsilon_0]$ can be represented as, respectively, $(h'_A, k'_A, \delta'_A, G'_A, \varepsilon'_A)$ - and $(h''_A, k''_A, \delta''_A, G''_A, \varepsilon''_A)$ -expansion, then the (h_A, k_A) -expansion $A(\varepsilon) = a_{h_A} \varepsilon^{h_A} + \dots + a_{k_A} \varepsilon^{k_A} + o_A(\varepsilon^{k_A})$, $\varepsilon \in (0, \varepsilon_0]$ given in Lemma 1 $_*$ is an $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion, with parameters G_A, δ_A and ε_A chosen in the following way consistent with the priority order described above:*

$$(\delta_A, G_A, \varepsilon_A) = \begin{cases} (\delta''_A, G''_A, \varepsilon''_A) & \text{if } k'_A < k''_A \text{ or } k'_A = k''_A, \\ & \delta'_A < \delta''_A, \\ (\delta'_A = \delta''_A, G'_A \wedge G''_A, \varepsilon'_A \wedge \varepsilon''_A) & \text{if } k'_A = k''_A, \delta'_A = \delta''_A, \\ (\delta'_A, G'_A, \varepsilon'_A) & \text{if } k'_A > k''_A \text{ or } k'_A = k''_A, \\ & \delta'_A > \delta''_A. \end{cases}$$

It is also useful to mention that a constant a can be interpreted as function $A(\varepsilon) \equiv a$. Thus, 0 can be represented, for any integer $-\infty < h \leq k < \infty$, as the $(h, k, \delta_{h,k}, G_{h,k}, \varepsilon_{h,k})$ -expansion, $0 = 0\varepsilon^h + \dots + 0\varepsilon^k + o(\varepsilon^k)$, with remainder $o(\varepsilon^k) \equiv 0$ and, thus, parameters $\delta_{h,k} = 1$, an arbitrary small $G_{h,k} \in (0, \infty)$, and $\varepsilon_{h,k} = \varepsilon_0$. Also, 1 can be represented, for any integer $0 \leq k < \infty$, as the $(0, k, \delta_k, G_k, \varepsilon_k)$ -expansion, $1 = 1 + 0\varepsilon + \dots + 0\varepsilon^k + o(\varepsilon^k)$, with remainder $o(\varepsilon^k) \equiv 0$ and, thus, parameters $\delta_k = 1$, an arbitrary small $G_k \in (0, \infty)$, and $\varepsilon_k = \varepsilon_0$.

Let us consider four Laurent asymptotic expansions, $A(\varepsilon) = a_{h_A} \varepsilon^{h_A} + \dots + a_{k_A} \varepsilon^{k_A} + o_A(\varepsilon^{k_A})$, $B(\varepsilon) = b_{h_B} \varepsilon^{h_B} + \dots + b_{k_B} \varepsilon^{k_B} + o_B(\varepsilon^{k_B})$, $C(\varepsilon) =$

$c_{h_C}\varepsilon^{h_C} + \dots + c_{k_C}\varepsilon^{k_C} + o_C(\varepsilon^{k_C})$, and $D(\varepsilon) = d_{h_D}\varepsilon^{h_D} + \dots + d_{k_D}\varepsilon^{k_D} + o_D(\varepsilon^{k_D})$ defined for $0 < \varepsilon \leq \varepsilon_0$, for some $0 < \varepsilon_0 \leq 1$.

The following Lemma presents operational rules for computing parameters of upper bounds for remainders of Laurent asymptotic expansions.

Lemma 2. *The above asymptotic expansions have the following operational rules for computing remainders:*

(i) *If $A(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion and c is a constant, then $C(\varepsilon) = cA(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a $(h_C, k_C, \delta_C, G_C, \varepsilon_C)$ -expansion with parameters h_C, k_C and coefficients $c_r, r = h_C, \dots, k_C$ given in proposition (i) of Lemma 2*, and parameters δ_C, G_C and ε_C given by the formulas:*

$$(a) \delta_C = \delta_A;$$

$$(b) G_C = |c|G_A;$$

$$(c) \varepsilon_C = \varepsilon_A.$$

(ii) *If $A(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion and $B(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a $(h_B, k_B, \delta_B, G_B, \varepsilon_B)$ -expansion, then $C(\varepsilon) = A(\varepsilon) + B(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a $(h_C, k_C, \delta_C, G_C, \varepsilon_C)$ -expansion with parameters h_C, k_C and coefficients $c_r, r = h_C, \dots, k_C$ given in proposition (ii) of Lemma 2*, and parameters δ_C, G_C and ε_C given by formulas:*

$$(a) \delta_C = \delta_A I(k_A < k_B) + (\delta_A \wedge \delta_B) I(k_A = k_B) + \delta_B I(k_B < k_A) \geq \delta_A \wedge \delta_B;$$

$$(b) G_C = G_A \varepsilon_C^{k_A + \delta_A - k_C - \delta_C} + \sum_{k_C < i \leq k_A} |a_i| \varepsilon_C^{i - k_C - \delta_C} + G_B \varepsilon_C^{k_B + \delta_B - k_C - \delta_C} + \sum_{k_C < j \leq k_B} |b_j| \varepsilon_C^{j - k_C - \delta_C};$$

$$(c) \varepsilon_C = \varepsilon_A \wedge \varepsilon_B.$$

(iii) *If $A(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion and $B(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a $(h_B, k_B, \delta_B, G_B, \varepsilon_B)$ -expansion, then $C(\varepsilon) = A(\varepsilon) \cdot B(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a $(h_C, k_C, \delta_C, G_C, \varepsilon_C)$ -expansion with parameters h_C, k_C and coefficients $c_r, r = h_C, \dots, k_C$ given in proposition (iii) of Lemma 2*, and parameters δ_C, G_C and ε_C given by formulas:*

$$(a) \delta_C = \delta_A I(k_A + h_B < k_B + h_A) + (\delta_A \wedge \delta_B) I(k_A + h_B = k_B + h_A) + \delta_B I(k_A + h_B > k_B + h_A) \geq \delta_A \wedge \delta_B;$$

$$(b) G_C = \sum_{k_C < i+j, h_A \leq i \leq k_A, h_B \leq j \leq k_B} |a_i| |b_j| \varepsilon_C^{i+j-k_C-\delta_C} + G_A \sum_{h_B \leq j \leq k_B} |b_j| \varepsilon_C^{j+k_A+\delta_A-k_C-\delta_C} + G_B \sum_{h_A \leq i \leq k_A} |a_i| \varepsilon_C^{i+k_B+\delta_B-k_C-\delta_C} + G_A G_B \varepsilon_C^{k_A+k_B+\delta_A+\delta_B-k_C-\delta_C};$$

$$(c) \varepsilon_C = \varepsilon_A \wedge \varepsilon_B.$$

(iv) *If $B(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a pivotal $(h_B, k_B, \delta_B, G_B, \varepsilon_B)$ -expansion, then there exists $\varepsilon_C \leq \varepsilon'_0 \leq \varepsilon_0$ such that $B(\varepsilon) \neq 0, \varepsilon \in (0, \varepsilon'_0]$, and $C(\varepsilon) = \frac{1}{B(\varepsilon)}, \varepsilon \in (0, \varepsilon'_0]$ is a pivotal $(h_C, k_C, \delta_C, G_C, \varepsilon_C)$ -expansion with parameters h_C, k_C and*

coefficients $c_r, r = h_C, \dots, k_C$ given in proposition (iv) of Lemma 2*, and parameters δ_C, G_C and ε_C given formulas:

(a) $\delta_C = \delta_B$;

(b) $G_C = (\frac{|b_{h_B}|}{2})^{-1} (\sum_{k_B - h_B < i+j, h_B \leq i \leq k_B, h_C \leq j \leq k_C} |b_i| |c_j| \varepsilon_C^{i+j-k_B+h_B-\delta_B} + G_B \sum_{h_C \leq j \leq k_C} |c_j| \varepsilon_C^{j+h_B})$;

(c) $\varepsilon_C = \varepsilon_B \wedge \tilde{\varepsilon}_B$, where $\tilde{\varepsilon}_B = (\frac{|b_{h_B}|}{2(\sum_{h_B < i \leq k_B} |b_i| \varepsilon_B^{i-h_B-\delta_B}) + G_B \varepsilon_B^{k_B-h_B}})^{\frac{1}{\delta_B}}$.

(v) If $A(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion, $B(\varepsilon), \varepsilon \in (0, \varepsilon_0]$ is a pivotal $(h_B, k_B, \delta_B, G_B, \varepsilon_B)$ -expansion, then there exists $\varepsilon_D \leq \varepsilon'_0 \leq \varepsilon_0$ such that $B(\varepsilon) \neq 0, \varepsilon \in (0, \varepsilon'_0]$, and $D(\varepsilon) = \frac{A(\varepsilon)}{B(\varepsilon)}$ is a $(h_D, k_D, \delta_D, G_D, \varepsilon_D)$ -expansion with parameters h_D, k_D and coefficients $d_r, r = h_D, \dots, k_D$ given in proposition (v) of Lemma 2*, and parameters $\delta_D, G_D, \varepsilon_D$ given by formulas:

(a) $\delta_D = \delta_A I(h_C + k_A < h_A + k_C) + (\delta_A \wedge \delta_C) I(h_C + k_A = h_A + k_C) + \delta_C I(h_A + k_C < h_C + k_A) \geq \delta_A \wedge \delta_B$;

(b) $G_D = \sum_{k_D < i+j, h_A \leq i \leq k_A, h_C \leq j \leq k_C} |a_i| |c_j| \varepsilon_D^{i+j-k_D-\delta_D} + G_A \sum_{h_C \leq j \leq k_C} |c_j| \varepsilon_D^{j+k_A+\delta_A-k_D-\delta_D} + G_C \sum_{h_A \leq i \leq k_A} |a_i| \varepsilon_D^{i+k_C+\delta_C-k_D-\delta_D} + G_A G_C \varepsilon_D^{k_A+k_C+\delta_A+\delta_C-k_D-\delta_D}$;

(c) $\varepsilon_D = \varepsilon_A \wedge \varepsilon_C$,

where coefficients $c_r, r = h_C, \dots, k_C$ and parameters $h_C, k_C, \delta_C, G_C, \varepsilon_C$ are given for the $(h_C, k_C, \delta_C, G_C, \varepsilon_C)$ -expansion of function $C(\varepsilon) = \frac{1}{B(\varepsilon)}$ in the above proposition (iv), or by formulas:

(d) $\delta_D = \delta_A I(k_A - h_B < k_B - 2h_B + h_A) + (\delta_A \wedge \delta_B) I(k_A - h_B = k_B - 2h_B + h_A) + \delta_B I(k_A - h_B > k_B - 2h_B + h_A) \geq \delta_A \wedge \delta_B$;

(e) $G_D = (\frac{|b_{h_B}|}{2})^{-1} (\sum_{k_A \wedge (h_A+k_B-h_B) < i+j, h_A \leq i \leq k_A, h_D \leq j \leq k_D} |a_i| |d_j| \varepsilon_D^{i+j-k_D-h_B-\delta_D} + \sum_{k_A \wedge (h_A+k_B-h_B) < i \leq k_A} |a_i| \varepsilon_D^{i-h_B-k_D-\delta_D} + G_A \varepsilon_D^{k_A \delta_A - h_B - k_D - \delta_D} + G_B \sum_{h_D \leq j \leq k_D} |d_j| \varepsilon_D^{j+k_B+\delta_B-h_B-k_D-\delta_D})$;

(f) $\varepsilon_D = \varepsilon_A \wedge \varepsilon_B \wedge \tilde{\varepsilon}_B$, where $\tilde{\varepsilon}_B$ is given in relations (c) of proposition (iv).

Remark 1. Coefficients $\varepsilon_B, \varepsilon_C, \varepsilon_D \in (0, 1]$ are taken to nonnegative powers in all terms penetrating the sums, which define parameters G_C, G_D and $\tilde{\varepsilon}_B$, in Lemma 2.

The following operational rules for computing remainders for multiple summation and multiplication of Laurent asymptotic expansions, used in what follows, are analogues of the corresponding summation and multiplication rules given in Lemma 2.

Lemma 3. Let $A_m(\varepsilon) = a_{h_{A_m}, m} \varepsilon^{h_{A_m}} + \dots + a_{k_{A_m}, m} \varepsilon^{k_{A_m}} + o(\varepsilon^{k_{A_m}}), \varepsilon \in$

$(0, \varepsilon_0]$ be a $(h_{A_m}, k_{A_m}, \delta_{A_m}, G_{A_m}, \varepsilon_{A_m})$ -expansion, for $m = 1, \dots, N$. In this case:

(i) $B_n(\varepsilon) = A_1(\varepsilon) + \dots + A_n(\varepsilon)$, $\varepsilon \in (0, \varepsilon_0]$ is, for every $n = 1, \dots, N$, a $(h_{B_n}, k_{B_n}, \delta_{B_n}, G_{B_n}, \varepsilon_{B_n})$ -expansion, with parameters h_{B_n}, k_{B_n} , $n = 1, \dots, N$ and coefficients $b_{h_{B_n}+l, n}$, $l = 0, \dots, k_{B_n} - h_{B_n}$, $n = 1, \dots, N$ given in proposition (i) of Lemma 3*, and parameters $G_{B_n}, \delta_{B_n}, \varepsilon_{B_n}$, $n = 1, \dots, N$ given by formulas:

(a) $\delta_{B_n} = \min_{m \in \mathbb{K}_n} \delta_{A_m} \geq \delta_N^* = \min_{1 \leq m \leq n} \delta_{A_m}$, where $\mathbb{K}_n = \{m : 1 \leq m \leq n\}$,

$$k_m = \min(k_1, \dots, k_n);$$

(b) $G_{B_n} = \sum_{1 \leq i \leq n} (G_{A_i} \varepsilon_{B_n}^{k_{A_i} + \delta_{A_i} - k_{B_n} - \delta_{B_n}} + \sum_{k_{B_n} < j \leq k_{A_i}} |a_{A_i, j}| \varepsilon_{B_n}^{j - k_{B_n} - \delta_{B_n}});$

(c) $\varepsilon_{B_n} = \min(\varepsilon_{A_1}, \dots, \varepsilon_{A_n})$.

(ii) $C_n(\varepsilon) = A_1(\varepsilon) \times \dots \times A_n(\varepsilon)$, $\varepsilon \in (0, \varepsilon_0]$ is, for $n = 1, \dots, N$, a $(h_{C_n}, k_{C_n}, \delta_{C_n}, G_{C_n}, \varepsilon_{C_n})$ -expansion with parameters h_{C_n}, k_{C_n} , $n = 1, \dots, N$ and coefficients $c_{h_{C_n}+l, n}$, $l = 0, \dots, k_{C_n} - h_{C_n}$, $n = 1, \dots, N$ given in proposition (ii) of Lemma 3*, and parameters $G_{C_n}, \delta_{C_n}, \varepsilon_{C_n}$, $n = 1, \dots, N$ given by formulas:

(a) $\delta_{C_n} = \min_{m \in \mathbb{L}_n} \delta_{A_m} \geq \delta_N^*$, where $\mathbb{L}_n = \{m : 1 \leq m \leq n, (k_{A_m} + \sum_{1 \leq r \leq n, r \neq m} h_{A_r}) = \min_{1 \leq l \leq n} (k_{A_l} + \sum_{1 \leq r \leq n, r \neq l} h_{A_r})\}$;

(b) $G_{C_n} = \sum_{k_{C_n} < l_1 + \dots + l_n, h_{A_i} \leq l_i \leq k_{A_i}, 1 \leq i \leq n} \prod_{1 \leq i \leq n} |a_{A_i, l_i}| \varepsilon_{C_n}^{l_1 + \dots + l_n - k_{C_n} - \delta_{C_n}} + \sum_{1 \leq j \leq n} \prod_{1 \leq i \leq n, i \neq j} (\sum_{h_{A_i} \leq l \leq k_{A_i}} |a_{A_i, l}| \varepsilon_{C_n}^l + G_{A_i} \varepsilon_{C_n}^{k_{A_i} + \delta_{A_i}}) G_{A_j} \varepsilon_{C_n}^{k_{A_j} + \delta_{A_j} - k_{C_n} - \delta_{C_n}};$

(c) $\varepsilon_{C_n} = \min_{1 \leq i \leq n} \varepsilon_{A_i}$.

(iii) Parameters $\delta_{C_n}, G_{C_n}, \varepsilon_{C_n}$, $n = 1, \dots, N$ in upper bounds for remainders in the asymptotic expansions for functions $B_n(\varepsilon) = A_1(\varepsilon) + \dots + A_n(\varepsilon)$, $n = 1, \dots, N$ and $C_n(\varepsilon) = A_1(\varepsilon) \times \dots \times A_n(\varepsilon)$, $n = 1, \dots, N$ are invariant with respect to any permutation, respectively, of summation and multiplication order in the above formulas.

The summation and multiplication rules for computing of upper bounds for remainders given in propositions (ii) and (iii) of Lemma 2 possess the communicative property, but do not possess the associative and distributal properties.

Lemma 2 let us get an effective low bound for parameter δ_A for any $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion $A(\varepsilon)$ obtained as the result of a finite sequence of operations (described in Lemma 2) performed over expansions from some finite set of such expansions.

The following lemma summarize these properties of Laurent asymptotic expansions with explicit upper bounds for remainders.

Lemma 4. *The summation and multiplication operations for Laurent asymptotic expansions defined in Lemma 2 possess the following algebraic properties, which should be understood as equalities for the corresponding parameters of upper bounds for their remainders:*

(i) *The functional identity, $C(\varepsilon) \equiv A(\varepsilon) + B(\varepsilon) \equiv B(\varepsilon) + A(\varepsilon)$, implies that $\delta_C = \delta_{A+B} = \delta_{B+A}$, $G_C = G_{A+B} = G_{B+A}$ and $\varepsilon_C = \varepsilon_{A+B} = \varepsilon_{B+A}$.*

(ii) *The functional identity, $C(\varepsilon) \equiv A(\varepsilon) \cdot B(\varepsilon) \equiv B(\varepsilon) \cdot A(\varepsilon)$, implies that $\delta_C = \delta_{A \cdot B} = \delta_{B \cdot A}$, $G_C = G_{A \cdot B} = G_{B \cdot A}$ and $\varepsilon_C = \varepsilon_{A \cdot B} = \varepsilon_{B \cdot A}$.*

(iii) *If $A(\varepsilon)$ is $(h_A, k_A, \delta_A, G_A, \varepsilon_A)$ -expansion obtained as the result of a finite sequence of operations (multiplication by a constant, summation, multiplication, and division) performed over $(h_{A_i}, k_{A_i}, \delta_{A_i}, G_{A_i}, \varepsilon_{A_i})$ -expansions $A_i(\varepsilon)$, $i = 1, \dots, N$, according the rules presented in Lemmas 2* and 2, then $\delta_A \geq \delta_N^* = \min_{1 \leq i \leq N} \delta_{A_i}$. This makes it possible to rewrite $A(\varepsilon)$ as the $(h_A, k_A, \delta_N^*, G_{A,N}^*, \varepsilon_A)$ -expansion, with parameter $G_{A,N}^* = G_A \varepsilon_A^{\delta_A - \delta_N^*}$.*

3. Asymptotic expansions for transition characteristics of non-linearly perturbed semi-Markov processes with reduced phase spaces

Let us recall the perturbed semi-Markov processes $\eta^{(\varepsilon)}(t)$, $t \geq 0$, with phase space $\mathbb{X} = \{1, \dots, N\}$ and transition probabilities $Q_{ij}^{(\varepsilon)}(t)$, $t \geq 0$, $i, j \in \mathbb{X}$, introduced in Part I of the paper. These processes depend on a perturbation parameter $\varepsilon \in (0, \varepsilon_0]$, for some $0 < \varepsilon_0 \leq 1$. We also recall transition probabilities of the corresponding embedded Markov chains, $p_{ij}(\varepsilon) = Q_{ij}^{(\varepsilon)}(\infty)$, $i, j \in \mathbb{X}$, and expectations of sojourn times $e_{ij}(\varepsilon) = \int_0^\infty t Q_{ij}^{(\varepsilon)}(dt)$, $i, j \in \mathbb{X}$.

We assume that condition **A**, introduced in Part I, holds for semi-Markov processes $\eta^{(\varepsilon)}(t)$. In particular, we recall the transition sets $\mathbb{Y}_i$, $i \in \mathbb{X}$ (which include states $j \in \mathbb{X}$ with non-zero probabilities $p_{ij}(\varepsilon)$ and guarantee ergodicity of the processes $\eta^{(\varepsilon)}(t)$) introduced in this condition.

However, we replace the perturbation condition **D** by the following stronger condition, in which the corresponding Taylor asymptotic expansions are given in the form with explicit upper bounds for remainders:

$$\mathbf{D}': p_{ij}(\varepsilon) = \sum_{l=l_{ij}^-}^{l_{ij}^+} a_{ij}[l] \varepsilon^l + o_{ij}(\varepsilon^{l_{ij}^+}), \varepsilon \in (0, \varepsilon_0], \text{ for } j \in \mathbb{Y}_i, i \in \mathbb{X}, \text{ where } (\mathbf{a})$$

$a_{ij}[l_{ij}^-] > 0$ and $0 \leq l_{ij}^- \leq l_{ij}^+ < \infty$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$; **(b)** $|o_{ij}(\varepsilon^{l_{ij}^+})| \leq G_{ij}\varepsilon^{l_{ij}^+ + \delta_{ij}}, 0 < \varepsilon \leq \varepsilon_{ij}$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$, where $0 < \delta_{ij} \leq 1, 0 < G_{ij} < \infty$ and $0 < \varepsilon_{ij} \leq \varepsilon_0$.

Also, we replace the perturbation condition **E** by the following stronger condition, in which the corresponding Laurent asymptotic expansions are given in the form with explicit upper bounds for remainders:

E': $e_{ij}(\varepsilon) = \sum_{l=m_{ij}^-}^{m_{ij}^+} b_{ij}[l]\varepsilon^l + \dot{o}_{ij}(\varepsilon^{m_{ij}^+}), \varepsilon \in (0, \varepsilon_0]$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$, where
(a) $b_{ij}[l_{ij}^-] > 0$ and $-\infty < m_{ij}^- \leq m_{ij}^+ < \infty$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$; **(b)**
 $|\dot{o}_{ij}(\varepsilon^{l_{ij}^+})| \leq \dot{G}_{ij}\varepsilon^{m_{ij}^+ + \dot{\delta}_{ij}}, 0 < \varepsilon \leq \dot{\varepsilon}_{ij}$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$, where $0 < \dot{\delta}_{ij} \leq 1, 0 < \dot{G}_{ij} < \infty$ and $0 < \dot{\varepsilon}_{ij} \leq \varepsilon_0$.

As was pointed out in Part I, condition **A** implies that sets $\mathbb{Y}_{rr}^+ = \mathbb{Y}_{rr} \setminus \{r\} \neq \emptyset, r \in \mathbb{X}$ and the non-absorption probability $\bar{p}_{rr}(\varepsilon) = 1 - p_{rr}(\varepsilon) \in (0, 1]$, for $r \in \mathbb{X}, \varepsilon \in (0, \varepsilon_0]$. This probability satisfy the following relation, for every $r \in \mathbb{X}, \varepsilon \in (0, \varepsilon_0]$,

$$\bar{p}_{rr}(\varepsilon) = 1 - p_{rr}(\varepsilon) = \sum_{j \in \mathbb{Y}_{rr}^+} p_{rj}(\varepsilon). \quad (2)$$

The above relation let us construct an algorithm for getting asymptotic expansions with explicit upper bounds for remainders, for non-absorption probabilities $\bar{p}_{rr}(\varepsilon)$.

Lemma 5. *Let conditions **A** and **D'** hold. Then, for every $r \in \mathbb{X}$, the pivotal $(\bar{l}_{rr}^-, \bar{l}_{rr}^+)$ -expansion for the non-absorption probability $\bar{p}_{rr}(\varepsilon)$ given in Lemma 8* is, also, a $(\bar{l}_{rr}^-, \bar{l}_{rr}^+, \bar{\delta}_{rr}, \bar{G}_{rr}, \bar{\varepsilon}_{rr})$ -expansion, with parameters $\bar{\delta}_{rr}, \bar{G}_{rr}$ and $\bar{\varepsilon}_{rr}$, which can be computed according the algorithm described below, in the proof of the lemma.*

Proof. Let $r \in \mathbb{Y}_r$. First, propositions **(i)** of Lemmas 3* and 3 (the multiple summation rule) should be applied to the sum $\sum_{j \in \mathbb{Y}_{rr}^+} p_{rj}(\varepsilon)$. Second, propositions **(i)** (the multiplication by constant -1) and **(ii)** (the summation with constant 1) of Lemmas 2* and 2 should be applied to the asymptotic expansion for probability $p_{rr}(\varepsilon)$ given in condition **D'**, in order to get the asymptotic expansion for function $1 - p_{rr}(\varepsilon)$. Third, Lemmas 1* and 1 should be applied to the asymptotic expansion for function $\bar{p}_{rr}(\varepsilon)$ given in two alternative forms by relation (2). This yields the corresponding pivotal the $(\bar{l}_{rr}^-, \bar{l}_{rr}^+)$ -expansion for probabilities $\bar{p}_{rr}(\varepsilon)$, given in Lemma 8*, and

proves that this expansion is a $(\bar{l}_{rr}^-, \bar{l}_{rr}^+, \bar{\delta}_{rr}, \bar{G}_{rr}, \bar{\varepsilon}_{rr})$ -expansion, with parameters computed in the process of realization of the above algorithm. The case $r \notin \mathbb{Y}_r$ is trivial, since, in this case, probability $\bar{p}_{rr}(\varepsilon) \equiv 1$. $\square$

Let us recall formula (19)_{*} for the transition probabilities ${}_r p_{ij}(\varepsilon)$, $i, j \in {}_r \mathbb{X} = \mathbb{X} \setminus \{r\}$ of the reduced embedded Markov chain ${}_r \eta_n^{(\varepsilon)}$, introduced in Part I,

$${}_r p_{ij}(\varepsilon) = p_{ij}(\varepsilon) + p_{ir}(\varepsilon) \frac{p_{rj}(\varepsilon)}{1 - p_{rr}(\varepsilon)}. \quad (3)$$

Let us introduce parameter,

$$\delta^\circ = \min_{j \in \mathbb{Y}_i, i \in \mathbb{X}} \delta_{ij}. \quad (4)$$

Obviously, inequalities $\delta_{ij} \geq \delta^\circ$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$ hold for parameters δ_{ij} penetrating upper bounds for the remainders of asymptotic expansions in condition **D'**.

Theorem 1. *Conditions **A** and **D'**, assumed to hold for the Markov chains $\eta_n^{(\varepsilon)}$, also hold for the reduced Markov chains ${}_r \eta_n^{(\varepsilon)}$, for every $r \in \mathbb{X}$. Also, for every $j \in {}_r \mathbb{Y}_i$, $i \in {}_r \mathbb{X}$, $r \in \mathbb{X}$, the pivotal $({}_r l_{ij}^-, {}_r l_{ij}^+)$ -expansion for the transition probability ${}_r p_{ij}(\varepsilon)$ given in Theorem 2_{*} is a $({}_r l_{ij}^-, {}_r l_{ij}^+, {}_r \delta_{ij}, {}_r G_{ij}, {}_r \varepsilon_{ij})$ -expansion penetrating condition **D'** for the Markov chains ${}_r \eta_n^{(\varepsilon)}$. Parameters ${}_r \delta_{ij}$, ${}_r G_{ij}$ and ${}_r \varepsilon_{ij}$ can be computed using the algorithm described below, in the proof of the theorem. The inequalities ${}_r \delta_{ij} \geq \delta^\circ$, $j \in {}_r \mathbb{Y}_i$, $i \in {}_r \mathbb{X}$, $r \in \mathbb{X}$ hold.*

Proof. Condition **A** holds for the Markov chains ${}_r \eta_n^{(\varepsilon)}$ by Lemma 6_{*}, with the same parameter ε_0 as for the Markov chains $\eta_n^{(\varepsilon)}$ and with the transition sets ${}_r \mathbb{Y}_i$, $i \in {}_r \mathbb{X}$ given by relation (20)_{*}.

Let us prove that condition **D'** holds for the Markov chains ${}_r \eta_n^{(\varepsilon)}$, with the same parameter ε_0 as for the Markov chains $\eta_n^{(\varepsilon)}$ and the transition sets ${}_r \mathbb{Y}_i$, $i \in {}_r \mathbb{X}$ given by relation (20)_{*}. Let $j, r \in \mathbb{Y}_i \cap \mathbb{Y}_r$. First, propositions **(v)** (the division rule) of Lemmas 2_{*} and 2 should be applied to the quotient $\frac{p_{rj}(\varepsilon)}{1 - p_{rr}(\varepsilon)}$. Second, propositions **(iii)** (the multiplication rule) of Lemmas 2_{*} and 2 should be applied to the product $p_{ir}(\varepsilon) \cdot \frac{p_{rj}(\varepsilon)}{1 - p_{rr}(\varepsilon)}$. Third, propositions **(ii)** (the summation rule) of Lemmas 2_{*} and 2 should be applied to sum ${}_r p_{ij}(\varepsilon) = p_{ij}(\varepsilon) + p_{ir}(\varepsilon) \cdot \frac{p_{rj}(\varepsilon)}{1 - p_{rr}(\varepsilon)}$. The asymptotic expansions for probabilities $p_{ir}(\varepsilon)$, $p_{rj}(\varepsilon)$, and $p_{ij}(\varepsilon)$, given in condition **D'**, and probability $1 - p_{rr}(\varepsilon)$, given in Lemmas 8_{*} and 5, should be used. This yields

the corresponding pivotal $({}_r l_{ij}^-, {}_r l_{ij}^+)$ -expansions for transition probabilities ${}_r p_{ij}(\varepsilon)$, $j \in {}_r \mathbb{Y}_i$, $i \in {}_r \mathbb{X}$, $r \in \mathbb{X}$, given in Theorem 2*, and proves that these expansions are $({}_r l_{ij}^-, {}_r l_{ij}^+, {}_r \delta_{ij}, {}_r \dot{G}_{ij}, {}_r \dot{\varepsilon}_{ij})$ -expansions, with parameters computed in the process of realization of the above algorithm. If $j \notin \mathbb{Y}_i$ then $p_{ij}(\varepsilon) \equiv 0$; if $j \notin \mathbb{Y}_r$ then $p_{rj}(\varepsilon) \equiv 0$; if $r \notin \mathbb{Y}_i$ then $p_{ir}(\varepsilon) \equiv 0$; if $r \notin \mathbb{Y}_r$ then $1 - p_{rr}(\varepsilon) \equiv 1$. In these cases, the above algorithm is readily simplified. Thus, condition **D'** holds for the reduced Markov chains ${}_r \eta_n^{(\varepsilon)}$.

Inequalities ${}_r \delta_{ij} \geq \delta^\circ$, $j \in {}_r \mathbb{Y}_i$, $i \in {}_r \mathbb{X}$, $r \in \mathbb{X}$ hold, by proposition (iii) of Lemma 4. $\square$

Let us recall formula (22)* for the expectations of sojourn times ${}_r e_{ij}(\varepsilon)$, for $i, j \in {}_r \mathbb{X} = \mathbb{X} \setminus \{r\}$ for the reduced semi-Markov process ${}_r \eta^{(\varepsilon)}(t)$, introduced in Part I,

$$\begin{aligned} {}_r e_{ij}(\varepsilon) &= e_{ij}(\varepsilon) + e_{ir}(\varepsilon) \frac{p_{rj}(\varepsilon)}{1 - p_{rr}(\varepsilon)} \\ &\quad + e_{rr}(\varepsilon) \frac{p_{ir}(\varepsilon)}{1 - p_{rr}(\varepsilon)} \frac{p_{rj}(\varepsilon)}{1 - p_{rr}(\varepsilon)} + e_{rj}(\varepsilon) \frac{p_{ir}(\varepsilon)}{1 - p_{rr}(\varepsilon)}. \end{aligned} \quad (5)$$

Let us introduce parameter,

$$\delta^* = \min_{j \in \mathbb{Y}_i, i \in \mathbb{X}} (\delta_{ij} \wedge \dot{\delta}_{ij}). \quad (6)$$

Obviously, inequalities $\delta_{ij}, \dot{\delta}_{ij} \geq \delta^*$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$ hold for parameters δ_{ij} and $\dot{\delta}_{ij}$ penetrating upper bounds for the remainders of asymptotic expansions in conditions **D'** and **E'**.

Theorem 2. *Conditions **A** – **C**, **D'** and **E'**, assumed to hold for the semi-Markov processes $\eta^{(\varepsilon)}(t)$, also hold for the reduced semi-Markov processes ${}_r \eta^{(\varepsilon)}(t)$, for every $r \in \mathbb{X}$. Also, for every $j \in {}_r \mathbb{Y}_i$, $i \in {}_r \mathbb{X}$, $r \in \mathbb{X}$, the pivotal $({}_r m_{ij}^-, {}_r m_{ij}^+)$ -expansion for expectation ${}_r e_{ij}(\varepsilon)$ given in Theorem 3* is a $({}_r m_{ij}^-, {}_r m_{ij}^+, {}_r \dot{\delta}_{ij}, {}_r \dot{G}_{ij}, {}_r \dot{\varepsilon}_{ij})$ -expansion penetrating condition **E'** for the semi-Markov processes ${}_r \eta^{(\varepsilon)}(t)$. Parameters ${}_r \dot{\delta}_{ij}$, ${}_r \dot{G}_{ij}$ and ${}_r \dot{\varepsilon}_{ij}$ can be computed using the algorithm described below, in the proof of the theorem. The inequalities ${}_r \dot{\delta}_{ij} \geq \delta^*$, $j \in {}_r \mathbb{Y}_i$, $i \in {}_r \mathbb{X}$, $r \in \mathbb{X}$ hold.*

Proof. Conditions **A** and **D'** hold for the semi-Markov processes ${}_r \eta^{(\varepsilon)}(t)$, respectively, by Lemma 6* and Theorem 1, with the same parameter ε_0 as for the semi-Markov processes $\eta^{(\varepsilon)}(t)$, and the transition sets ${}_r \mathbb{Y}_i$, $i \in {}_r \mathbb{X}$ given by relation (20)*. Also conditions **B** and **C** hold for processes ${}_r \eta^{(\varepsilon)}(t)$, by Lemma 7*.

Let us prove that condition **E'** holds for the semi-Markov processes ${}_r\eta^{(\varepsilon)}(t)$, with the same parameter ε_0 and the transition sets ${}_r\mathbb{Y}_i, i \in {}_r\mathbb{X}$ given by relation (20)*. Let $j, r \in \mathbb{Y}_i \cap \mathbb{Y}_r$. First, propositions **(v)** (the division rule) of Lemmas 2* and 2 should be applied to the quotients $\frac{p_{rj}(\varepsilon)}{1-p_{rr}(\varepsilon)}$ and $\frac{p_{ir}(\varepsilon)}{1-p_{rr}(\varepsilon)}$. Second, propositions **(iii)** (the multiplication rule) of Lemmas 2* and 2 should be applied to the products $e_{ir}(\varepsilon) \cdot \frac{p_{rj}(\varepsilon)}{1-p_{rr}(\varepsilon)}$ and $e_{rj}(\varepsilon) \cdot \frac{p_{ir}(\varepsilon)}{1-p_{rr}(\varepsilon)}$ and propositions **(ii)** (the multiple multiplication rule) of Lemmas 3* and 3 to the product $e_{rr}(\varepsilon) \cdot \frac{p_{ir}(\varepsilon)}{1-p_{rr}(\varepsilon)} \cdot \frac{p_{rj}(\varepsilon)}{1-p_{rr}(\varepsilon)}$. Third, propositions **(i)** (the multiple summation rule) of Lemmas 3* and 3 should be applied to sum ${}_re_{ij}(\varepsilon) = e_{ij}(\varepsilon) + e_{ir}(\varepsilon) \cdot \frac{p_{rj}(\varepsilon)}{1-p_{rr}(\varepsilon)} + e_{rr}(\varepsilon) \cdot \frac{p_{ir}(\varepsilon)}{1-p_{rr}(\varepsilon)} \cdot \frac{p_{rj}(\varepsilon)}{1-p_{rr}(\varepsilon)} + e_{rj}(\varepsilon) \cdot \frac{p_{ir}(\varepsilon)}{1-p_{rr}(\varepsilon)}$. The asymptotic expansions for probabilities $p_{ir}(\varepsilon), p_{ir}(\varepsilon)$ and $p_{ij}(\varepsilon)$, given in condition **D'**, probability $1-p_{rr}(\varepsilon)$, given in Lemmas 8* and 5, and expectations $e_{ij}(\varepsilon), e_{ir}(\varepsilon), e_{rr}(\varepsilon)$ and $e_{rj}(\varepsilon)$, given in condition **E'**, should be used. This, first, yields the corresponding pivotal $({}_rm_{ij}^-, {}_rm_{ij}^+)$ -expansions for expectations of sojourn times ${}_re_{ij}(\varepsilon), j \in {}_r\mathbb{Y}_i, i \in {}_r\mathbb{X}, r \in \mathbb{X}$, s given in Theorem 3*, and, second, proves that these expansions are $({}_rm_{ij}^-, {}_rm_{ij}^+, {}_r\dot{\delta}_{ij}, {}_r\dot{G}_{ij}, {}_r\dot{e}_{ij})$ -expansions, with parameters computed in the process of realization of the above algorithm. If $j \notin \mathbb{Y}_i$ then $p_{ij}(\varepsilon) \equiv 0$ and $e_{ij}(\varepsilon) \equiv 0$; if $j \notin \mathbb{Y}_r$ then $p_{rj}(\varepsilon) \equiv 0$ and $e_{rj}(\varepsilon) \equiv 0$; if $r \notin \mathbb{Y}_i$ then $p_{ir}(\varepsilon) \equiv 0$ and $e_{ir}(\varepsilon) \equiv 0$; if $r \notin \mathbb{Y}_r$ then $1-p_{rr}(\varepsilon) \equiv 1$ and $e_{rr}(\varepsilon) \equiv 0$. In these cases, the above algorithm is readily simplified. Thus, condition **E'** holds for the reduced semi-Markov processes ${}_r\eta^{(\varepsilon)}(t)$.

Inequalities ${}_r\dot{\delta}_{ij} \geq \delta^*, j \in {}_r\mathbb{Y}_i, i \in {}_r\mathbb{X}, r \in \mathbb{X}$ hold, by proposition **(iii)** of Lemma 4. $\square$

It worth to note that, despite bulky forms, formulas for parameters of upper bounds for remainders, in the asymptotic expansions given in Lemma 5 and Theorems 1 and 2, are computationally effective.

4. Asymptotic expansions for expected hitting times with explicit upper bounds for remainders

As in Part I, let $\bar{r}_{i,N} = \langle r_{i,1}, \dots, r_{i,N} \rangle = \langle r_{i,1}, \dots, r_{i,N-1}, i \rangle$ be a permutation of the sequence $\langle 1, \dots, N \rangle$ such that $r_{i,N} = i$, and let $\bar{r}_{i,n} = \langle r_{i,1}, \dots, r_{i,n} \rangle, n = 1, \dots, N$ be the corresponding chain of growing sequences of states from space $\mathbb{X}$.

Theorem 3. *Let conditions **A** – **C**, **D'** and **E'** hold for the semi-Markov*

processes $\eta^{(\varepsilon)}(t)$. Then, for every $i \in \mathbb{X}$, the pivotal (M_{ii}^-, M_{ii}^+) -expansion for the expectation of hitting time $E_{ii}(\varepsilon)$, given in Theorem 4* and obtained as the result of sequential exclusion of states $r_{i,1}, \dots, r_{i,N-1}$ from the phase space $\mathbb{X}$ of the processes $\eta^{(\varepsilon)}(t)$, is a $(M_{ii}^-, M_{ii}^+, \bar{r}_{i,N-1} \dot{\delta}_{ii}, \bar{r}_{i,N-1} \dot{G}_{ii}, \bar{r}_{i,N-1} \dot{\varepsilon}_{ii})$ -expansion. Parameters $\bar{r}_{i,N-1} \dot{\delta}_{ii}$, $\bar{r}_{i,N-1} \dot{G}_{ii}$ and $\bar{r}_{i,N-1} \dot{\varepsilon}_{ii}$ can be computed using the algorithm described below, in the proof of the theorem. Also, inequality $\bar{r}_{i,N-1} \dot{\delta}_{ii} \geq \delta^*$ holds making it possible to rewrite function $E_{ii}(\varepsilon)$ as the pivotal $(M_{ii}^-, M_{ii}^+, \delta^*, \bar{r}_{i,N-1} \dot{G}_{ii}^*, \bar{r}_{i,N-1} \dot{\varepsilon}_{ii})$ -expansion, with parameter $\bar{r}_{i,N-1} \dot{G}_{ii}^* = \bar{r}_{i,N-1} \dot{G}_{ii} \cdot (\bar{r}_{i,N-1} \dot{\varepsilon}_{ii})^{(\bar{r}_{i,N-1} \dot{\delta}_{ii} - \delta^*)}$.

Proof. Let us assume that $p_i^{(\varepsilon)} = 1$. Denote as $\bar{r}_{i,0} \eta^{(\varepsilon)}(t) = \eta^{(\varepsilon)}(t)$ the initial semi-Markov process. Let us exclude state $r_{i,1}$ from the phase space of semi-Markov process $\bar{r}_{i,0} \eta^{(\varepsilon)}(t)$ using the time-space screening procedure described in Section 5*. Let $\bar{r}_{i,1} \eta^{(\varepsilon)}(t)$ be the corresponding reduced semi-Markov process. The above procedure can be repeated. The state $r_{i,2}$ can be excluded from the phase space of the semi-Markov process $\bar{r}_{i,1} \eta^{(\varepsilon)}(t)$. Let $\bar{r}_{i,2} \eta^{(\varepsilon)}(t)$ be the corresponding reduced semi-Markov process. By continuing the above procedure for states $r_{i,3}, \dots, r_{i,n}$, we construct the reduced semi-Markov process $\bar{r}_{i,n} \eta^{(\varepsilon)}(t)$.

The process $\bar{r}_{i,n} \eta^{(\varepsilon)}(t)$ has the phase space $\bar{r}_{i,n} \mathbb{X} = \mathbb{X} \setminus \{r_{i,1}, r_{i,2}, \dots, r_{i,n}\}$. The transition probabilities $\bar{r}_{i,n} p_{i'j'}(\varepsilon)$, $i', j' \in \bar{r}_{i,n} \mathbb{X}$ and the expectations of sojourn times $\bar{r}_{i,n} e_{i'j'}(\varepsilon)$, $i', j' \in \bar{r}_{i,n} \mathbb{X}$ are determined for the process $\bar{r}_{i,n} \eta^{(\varepsilon)}(t)$ by the transition probabilities and the expectations of sojourn times for the process $\bar{r}_{i,n-1} \eta^{(\varepsilon)}(t)$, via relations (3) and (5).

By Theorem 1*, the expectation of hitting time $E_{i'j'}(\varepsilon)$ coincides for the semi-Markov processes $\bar{r}_{i,0} \eta^{(\varepsilon)}(t)$, $\bar{r}_{i,1} \eta^{(\varepsilon)}(t)$, $\dots$, $\bar{r}_{i,n} \eta^{(\varepsilon)}(t)$, for every $i', j' \in \bar{r}_{i,n} \mathbb{X}$.

By Theorems 2*, 3*, 1 and 2, the semi-Markov processes $\bar{r}_{i,n} \eta^{(\varepsilon)}(t)$ satisfy conditions **A** – **C**, **D'** and **E'**. The transition sets $\bar{r}_{i,n} \mathbb{Y}_{i'}, i' \in \bar{r}_{i,n} \mathbb{X}$, for the process $\bar{r}_{i,n} \eta^{(\varepsilon)}(t)$, are determined by the transition sets $\bar{r}_{i,n-1} \mathbb{Y}_{i'}, i' \in \bar{r}_{i,n-1} \mathbb{X}$, for the process $\bar{r}_{i,n-1} \eta^{(\varepsilon)}(t)$, via relation (20)*. For every $j' \in \bar{r}_{i,n} \mathbb{Y}_{i'}, i' \in \bar{r}_{i,n} \mathbb{X}$, the pivotal $(\bar{r}_{i,n} l_{i'j'}^-, \bar{r}_{i,n} l_{i'j'}^+)$ -expansion for transition probability $\bar{r}_{i,n} p_{i'j'}(\varepsilon)$, given in Theorem 2*, is, by Theorem 1, a $(\bar{r}_{i,n} l_{i'j'}^-, \bar{r}_{i,n} l_{i'j'}^+, \bar{r}_{i,n} \delta_{i'j'}, \bar{r}_{i,n} G_{i'j'}, \bar{r}_{i,n} \varepsilon_{i'j'})$ -expansion, with parameters $\bar{r}_{i,n} \delta_{i'j'}$, $\bar{r}_{i,n} G_{i'j'}$ and $\bar{r}_{i,n} \varepsilon_{i'j'}$ given in this theorem. Analogously, for every $j' \in \bar{r}_{i,n} \mathbb{Y}_{i'}, i' \in \bar{r}_{i,n} \mathbb{X}$, the pivotal $(\bar{r}_{i,n} m_{i'j'}^-, \bar{r}_{i,n} m_{i'j'}^+)$ -expansion for expectation $\bar{r}_{i,n} e_{i'j'}(\varepsilon)$, given in Theorem 3*, is, by Theorem 2, a $(\bar{r}_{i,n} m_{i'j'}^-, \bar{r}_{i,n} m_{i'j'}^+, \bar{r}_{i,n} \dot{\delta}_{i'j'}, \bar{r}_{i,n} \dot{G}_{i'j'}, \bar{r}_{i,n} \dot{\varepsilon}_{i'j'})$ -expansion, with pa-

rameters $\bar{r}_{i,n} \dot{\delta}_{i'j'}$, $\bar{r}_{i,n} \dot{G}_{i'j'}$ and $\bar{r}_{i,n} \dot{\varepsilon}_{i'j'}$ given in this theorem. Also, by Theorem 2, the inequalities $\bar{r}_{i,n} \dot{\delta}_{i'j'} \geq \delta^*$, $j' \in \bar{r}_{i,n} \mathbb{Y}_{i'}$, $i' \in \bar{r}_{i,n} \mathbb{X}$ hold.

The algorithm described above has a recurrent form and should be realized sequentially for the reduced semi-Markov processes $\bar{r}_{i,1} \eta^{(\varepsilon)}(t), \dots, \bar{r}_{i,n} \eta^{(\varepsilon)}(t)$ starting from the initial semi-Markov process $\bar{r}_{i,0} \eta^{(\varepsilon)}(t)$.

Let us take $n = N - 1$. The semi-Markov process $\bar{r}_{i,N-1} \eta^{(\varepsilon)}(t)$ has the phase space $\bar{r}_{i,N-1} \mathbb{X} = \mathbb{X} \setminus \{r_{i,1}, r_{i,2}, \dots, r_{i,N-1}\} = \{i\}$, which is a one-state set. The process $\bar{r}_{i,N-1} \eta^{(\varepsilon)}(t)$ returns to state i after every jump. Its transition probability $\bar{r}_{i,N-1} p_{ii}(\varepsilon) = 1$, and the expectation of hitting time, $E_{ii}(\varepsilon) = \bar{r}_{i,N-1} e_{ii}(\varepsilon)$. This equality and Theorem 4* yield, for every $i \in \mathbb{X}$, the pivotal (M_{ii}^-, M_{ii}^+) -expansion for expectation $E_{ii}(\varepsilon)$, which is invariant with respect to any permutation $\bar{r}'_{i,N-1} = \langle r'_{i,1}, \dots, r'_{i,N-1} \rangle$ of the sequence $\bar{r}_{i,N-1} = \langle r_{i,1}, \dots, r_{i,N-1} \rangle$. This invariance also implies that parameters $\bar{r}_{i,N-1} m_{ii}^\pm = M_{ii}^\pm$ do not depend on the choice of sequence $\bar{r}_{i,N-1}$. The above (M_{ii}^-, M_{ii}^+) -expansion for $E_{ii}(\varepsilon) = \bar{r}_{i,N-1} e_{ii}(\varepsilon)$ is a $(M_{ii}^-, M_{ii}^+, \bar{r}_{i,N-1} \dot{\delta}_{ii}, \bar{r}_{i,N-1} \dot{G}_{ii}, \bar{r}_{i,N-1} \dot{\varepsilon}_{ii})$ -expansion.

The above algorithm can be realized for any sequence $\bar{r}_{i,N-1} = \langle r_{i,1}, \dots, r_{i,N-1} \rangle$, but the invariance of explicit upper bounds for remainders, with respect to permutations $\bar{r}_{i,N} = \langle r_{i,1}, \dots, r_{i,N-1}, i \rangle$ of sequence $\langle 1, \dots, N \rangle$, can not be guaranteed.

However, the inequality $\bar{r}_{i,N-1} \dot{\delta}_{ii} \geq \delta^*$ holds, for any sequence $\bar{r}_{i,N-1} = \langle r_{i,1}, \dots, r_{i,N-1} \rangle$, by Theorem 2.

The algorithm described above can be repeated, for every $i \in \mathbb{X}$. $\square$

It is worth to note that the algorithms based on sequential exclusion of states from the phase space of perturbed semi-Markov processes make it possible to get Laurent asymptotic expansions (without and with explicit upper bounds for remainders) for expected hitting times, for nonlinearly perturbed semi-Markov processes. Such asymptotic results have their own important value.

Let $\bar{r}_{i,j,N} = \langle r_{i,j,1}, \dots, r_{i,j,N} \rangle = \langle r_{i,j,1}, \dots, r_{i,j,N-2}, i, j \rangle$ be a permutation of the sequence $\langle 1, \dots, N \rangle$ such that $r_{i,j,N-1} = i$, $r_{i,j,N} = j$, and let $\bar{r}_{i,j,n} = \langle r_{i,j,1}, \dots, r_{i,j,n} \rangle$, $n = 1, \dots, N$ be the corresponding chain of growing sequences of states from space $\mathbb{X}$.

By applying the algorithm of sequential phase space reduction described in Theorem 4 to the above sequence of states $\bar{r}_{i,j,N-2}$, we construct the reduced semi-Markov process $\bar{r}_{i,j,N-2} \eta^{(\varepsilon)}(t)$. This process has the phase space $\bar{r}_{i,j,N-2} \mathbb{X} = \mathbb{X}_{ij} = \{i, j\}$, which is a two-states set. The transition probabili-

ties of the embedded Markov chain $\bar{r}_{i,j,N-2} p_{i'j'}(\varepsilon) = p_{ij,i'j'}(\varepsilon)$, $i', j' \in \mathbb{X}_{ij}$, the expectations of sojourn times $\bar{r}_{i,j,N-2} e_{i'j'}(\varepsilon) = e_{ij,i'j'}(\varepsilon)$, $i', j' \in \mathbb{X}_{ij}$, and the transition sets $\bar{r}_{i,j,N-2} \mathbb{Y}_{i'} = \mathbb{Y}_{ij,i'}$, $i' \in \mathbb{X}_{ij}$ can be found using the recurrent algorithm described in Theorem 4*. These probabilities, expectations and transition sets are invariant to any permutation $\bar{r}'_{i,j,N-2}$ of sequence $\bar{r}_{i,j,N-2}$. This legitimates the above alternative simplified notations.

Theorem 4* let us construct the pivotal $(l_{ij,i'j'}^-, l_{ij,i'j'}^+)$ -expansions for the transition probabilities $p_{ij,i'j'}(\varepsilon)$, $j' \in \mathbb{Y}_{ij,i'}$, $i' \in \mathbb{X}_{ij}$ and the pivotal $(m_{ij,i'j'}^-, m_{ij,i'j'}^+)$ -expansions for the expectations of hitting times $e_{ij,i'j'}(\varepsilon)$, $j' \in \mathbb{Y}_{ij,i'}$, $i' \in \mathbb{X}_{ij}$, using the recurrent algorithm based on sequential exclusion states $r_{i,j,1}, \dots, r_{i,j,N-2}$ from the phase space $\mathbb{X}$. These expansions are invariant to any permutation $\bar{r}'_{i,j,N-2}$ of sequence $\bar{r}_{i,j,N-2}$.

According Theorem 3, the above $(l_{ij,i'j'}^-, l_{ij,i'j'}^+)$ -expansions for transition probabilities are $(l_{ij,i'j'}^-, l_{ij,i'j'}^+, \bar{r}_{i,j,N-2} \delta_{i'j'}, \bar{r}_{i,j,N-2} G_{i'j'}, \bar{r}_{i,j,N-2} \varepsilon_{i'j'})$ -expansions and the above $(m_{ij,i'j'}^-, m_{ij,i'j'}^+)$ -expansions for expectations of sojourn times are $(m_{ij,i'j'}^-, m_{ij,i'j'}^+, \bar{r}_{i,j,N-2} \dot{\delta}_{i'j'}, \bar{r}_{i,j,N-2} \dot{G}_{i'j'}, \bar{r}_{i,j,N-2} \dot{\varepsilon}_{i'j'})$ -expansions. This theorem let us also compute parameters $\bar{r}_{i,j,N-2} \delta_{i'j'}, \bar{r}_{i,j,N-2} G_{i'j'}, \bar{r}_{i,j,N-2} \varepsilon_{i'j'}$ and $\bar{r}_{i,j,N-2} \dot{\delta}_{i'j'}, \bar{r}_{i,j,N-2} \dot{G}_{i'j'}, \bar{r}_{i,j,N-2} \dot{\varepsilon}_{i'j'}$ of upper bounds for the corresponding remainders.

By Theorem 1*, the expectation of hitting time $E_{i'j'}(\varepsilon)$ coincides for the initial semi-Markov processes $\eta^{(\varepsilon)}(t)$ and the reduced semi-Markov process $\bar{r}_{i,j,N-2} \eta^{(\varepsilon)}(t)$, for every $i', j' \in \mathbb{X}_{ij}$. This obviously implies that these expectations are also invariant to any permutation $\bar{r}'_{i,j,N-2}$ of sequence $\bar{r}_{i,j,N-2}$.

It is easy to write down the formulas for the above expectations, for the two-states semi-Markov process $\bar{r}_{i,j,N-2} \eta^{(\varepsilon)}(t)$. These formulas are, $E_{i'j'}(\varepsilon) = e_{ij,i'j'}(\varepsilon) \frac{1}{1-p_{ij,i'j'}(\varepsilon)}$, $E_{j'j'}(\varepsilon) = e_{ij,j'j'}(\varepsilon) + e_{ij,j'j'}(\varepsilon) \frac{p_{ij,j'j'}(\varepsilon)}{1-p_{ij,i'j'}(\varepsilon)}$, where $e_{ij,i'j'}(\varepsilon) = e_{ij,i'j'}(\varepsilon) + e_{ij,i'j'}(\varepsilon)$, $i', j' \in \mathbb{X}_{ij}$, $i' \neq j'$.

Under the assumption that conditions of Theorem 4* hold, the operational rules given in Lemma 2* can be applied to functions $E_{i'j'}(\varepsilon)$, $i', j' \in \mathbb{X}_{ij}$, in order to get the corresponding $(M_{i'j'}^-, M_{i'j'}^+)$ -expansions. These expansions are invariant to any permutation $\bar{r}'_{i,j,N-2}$ of sequence $\bar{r}_{i,j,N-2}$, used in the corresponding recurrent algorithm based on sequential exclusion states $r_{i,j,1}, \dots, r_{i,j,N-2}$ from the phase space $\mathbb{X}$.

Finally, under the assumption that conditions of Theorem 3 hold, the operational rules given in Lemma 2 can be applied, in order to prove that the above $(M_{i'j'}^-, M_{i'j'}^+)$ -expansions are $(M_{i'j'}^-, M_{i'j'}^+, \bar{r}_{i,j,N-2} \dot{\delta}_{i'j'}, \bar{r}_{i,j,N-2} \dot{G}_{i'j'},$

$\bar{r}_{i,j,N-2}\dot{\varepsilon}_{i'j'}$)-expansions, and to compute parameters $\bar{r}_{i,j,N-2}\dot{\delta}_{i'j'}$, $\bar{r}_{i,j,N-2}\dot{G}_{i'j'}$ and $\bar{r}_{i,j,N-2}\dot{\varepsilon}_{i'j'}$ of upper bounds for the corresponding remainders. Also, by Lemma 4, the inequality $\bar{r}_{i,j,N-2}\dot{\delta}_{i'j'} \geq \delta^*$ holds, for every $i', j' \in \mathbb{X}_{ij}$, sequence $\bar{r}_{i,j,N-2}$, and $i, j \in \mathbb{X}$.

5. Asymptotic expansions for stationary distributions with explicit upper bounds for remainders

Let us recall the pivotal (n_i^-, n_i^+) -expansion for stationary probability $\pi_i(\varepsilon)$ of nonlinearly perturbed semi-Markov process $\eta^{(\varepsilon)}(t)$ given, under conditions **A** – **E**, in Theorem 5*. This asymptotic expansion has the following form, for $i \in \mathbb{X}$,

$$\pi_i(\varepsilon) = \sum_{l=n_i^-}^{n_i^+} c_i[l]\varepsilon^l + o_i(\varepsilon^{n_i^+}), \quad \varepsilon \in (0, \varepsilon_0]. \quad (7)$$

According Theorem 5*, the above asymptotic expansion is invariant with respect to the choice of sequence states $\bar{r}_{i,N-1} = (r_{i,1}, \dots, r_{i,N-1})$ used in the corresponding algorithm, for every $i \in \mathbb{X}$.

The following theorem is the main new result in Part II of this paper.

Theorem 4. *Let conditions **A** – **C**, **D'** and **E'** hold for the semi-Markov processes $\eta^{(\varepsilon)}(t)$. Then, for every $i \in \mathbb{X}$, the pivotal (n_i^-, n_i^+) -expansion (7) for the stationary probability $\pi_i(\varepsilon)$, given in Theorem 5* and obtained as the result of sequential exclusion of states $r_{i,1}, \dots, r_{i,N-1}$ from the phase space $\mathbb{X}$ of the processes $\eta^{(\varepsilon)}(t)$, is a $(n_i^-, n_i^+, \bar{r}_{i,N-1}\delta_i, \bar{r}_{i,N-1}G_i, \bar{r}_{i,N-1}\varepsilon_i)$ -expansion. Parameters $\bar{r}_{i,N-1}\delta_i$, $\bar{r}_{i,N-1}G_i$ and $\bar{r}_{i,N-1}\varepsilon_i$ can be computed using the algorithm described below, in the proof of the theorem. Also, inequality $\bar{r}_{i,N-1}\delta_i^* \geq \delta^*$ holds making it possible to rewrite function $\pi_i(\varepsilon)$ as the pivotal $(n_i^-, n_i^+, \delta^*, \bar{r}_{i,N-1}G_i^*, \bar{r}_{i,N-1}\varepsilon_i)$ -expansion, with parameter $\bar{r}_{i,N-1}G_i^* = \bar{r}_{i,N-1}G_i \cdot (\bar{r}_{i,N-1}\varepsilon_i)^{(\bar{r}_{i,N-1}\delta_i - \delta^*)}$.*

Proof. Let us choose an arbitrary state $i \in \mathbb{X}$. First, proposition (i) (the multiple summation rule) of Lemmas 3* and 3 should be applied to the pivotal (m_i^-, m_i^+) -expansion for the expectation $e_i(\varepsilon) = \sum_{j \in \mathbb{Y}_i} e_{ij}(\varepsilon)$ given by relation (29)*, in the proof of Theorem 5*. This yields a $(m_i^-, m_i^+, \dot{\delta}_i, \dot{G}_i, \dot{\varepsilon}_i)$ -expansion for the expectation $e_i(\varepsilon)$, with the corresponding parameters $\dot{\delta}_i, \dot{G}_i$ and $\dot{\varepsilon}_i^*$. Second, the propositions (v) (the division rule) of Lemmas 2* and

2 should be applied to the quotient $\pi_i(\varepsilon) = \frac{e_i(\varepsilon)}{E_{ii}(\varepsilon)}$. The $(m_i^-, m_i^+, \dot{\delta}_i, \dot{G}_i, \dot{\varepsilon}_i)$ -expansion for the expectation $e_i(\varepsilon)$ and the $(M_{ii}^-, M_{ii}^+, \bar{r}_{i,N-1}\dot{\delta}_{ii}, \bar{r}_{i,N-1}\dot{G}_{ii}, \bar{r}_{i,N-1}\dot{\varepsilon}_{ii})$ -expansion for the expectation of hitting time $E_{ii}(\varepsilon)$, given in Theorems 4* and 3, should be used. This yields the corresponding pivotal (n_i^-, n_i^+) -expansion for stationary probability $\pi_i(\varepsilon)$, given in Theorem 5*, and proves that this expansion is a $(n_i^-, n_i^+, \bar{r}_{i,N-1}\delta_i, \bar{r}_{i,N-1}G_i, \bar{r}_{i,N-1}\varepsilon_i)$ -expansion, with parameters computed in the process of realization of the above algorithm. Inequality $\bar{r}_{i,N-1}\delta_i \geq \delta^*$ holds, for every sequence $\bar{r}_{i,N-1}$, by proposition (iii) of Lemma 4. $\square$

The explicit upper bounds for remainders in the asymptotic expansions given in Theorem 4 have a clear and informative power-type form. An useful property of these upper bounds is that they are uniform with respect to the perturbation parameter. The recurrent algorithm for finding these upper bounds is computationally effective.

Unfortunately, the summation and multiplication operational rules for computing power-type upper bounds for remainders possess commutative but do not possess associative and distributive properties. This causes dependence of the resulting upper bounds for remainders in the asymptotic expansions for stationary probabilities $\pi_i(\varepsilon)$, $i \in \mathbb{X}$ on a choice of the corresponding sequences of states $\bar{r}_{i,N-1} = \langle r_{i,1}, \dots, r_{i,N-1} \rangle$, $i \in \mathbb{X}$ used in the above algorithm. This rises two open questions, the first one, about possible alternative forms for remainders possessing the desirable algebraic properties mentioned above, and, the second one, about an optimal choice of sequences of states $\bar{r}_{i,N-1}$, $i \in \mathbb{X}$.

In conclusion, we would like to mention some prospective directions for future research studies.

The method of sequential reduction of phase space presented in the paper can be applied for getting asymptotic expansions for high order power and exponential moments of hitting times, for nonlinearly perturbed semi-Markov processes. This is an interesting problem, which has its own important theoretical and applied values.

We are quite sure that a combination of results in the above direction with the methods of asymptotic analysis for nonlinearly perturbed regenerative processes developed and thoroughly presented in Gyllenberg and Silvestrov (2008) will make it possible to expand results from this book, related to asymptotic expansions for stationary and more general quasi-stationary distributions as well as other characteristics for nonlinearly perturbed semi-

Markov processes with absorption, to nonlinearly perturbed semi-Markov processes with an arbitrary asymptotic communicative structure of phase spaces.

The problems of aggregation of steps in the time-space screening procedures for semi-Markov processes, tracing pivotal orders for different groups of states as well as getting explicit formulas, for coefficients and parameters of upper bounds for remainders in the corresponding asymptotic expansions for stationary distributions and moments of hitting times, do require additional studies. It can be expected that such formulas can be obtained, for example, for nonlinearly perturbed birth-death-type semi-Markov processes, for which the proposed algorithms of phase space reduction preserve the birth-death structure for reduced semi-Markov processes.

Applications to control and queuing systems, information networks, epidemic models and models of mathematical genetics and population dynamics, analogous to those presented in the books cited in the introduction, also create a prospective area for future research based on the asymptotic results obtained in the present paper.

Appendix A: Operational rules for Laurent asymptotic expansions

Let us give short proofs of Lemmas $1_* - 4_*$ and $1 - 4$ omitting some known or obvious details.

A.1. The formulas given in Lemmas 1_* and 1 are quite obvious.

A.2. The same relates to formulas in propositions **(i)** (the multiplication by a constant rule) of Lemmas 2_* and 2 .

Proposition **(ii)** (the summation rules) of Lemmas 2_* and 2 can be obtained by simple accumulation of coefficients for different powers of ε and terms accumulated in the corresponding remainders, and, then, by using obvious upper bounds for absolute values of sums of terms accumulated in the corresponding remainders.

Proposition **(iii)** (the multiplication rule) of Lemma 2_* can be proved by multiplication of the corresponding asymptotic expansions $A(\varepsilon)$ and $B(\varepsilon)$ and accumulation of coefficients for powers ε^l for $l = h_C, \dots, k_C$ in their product,

$$\begin{aligned} C(\varepsilon) &= A(\varepsilon)B(\varepsilon) \\ &= (a_{h_A}\varepsilon^{h_A} + \dots + a_{k_A}\varepsilon^{k_A} + o_A(\varepsilon^{k_A})) \\ &\quad \times (b_{h_B}\varepsilon^{h_B} + \dots + b_{k_B}\varepsilon^{k_B} + o_B(\varepsilon^{k_B})) \end{aligned}$$

$$\begin{aligned}
&= \sum_{h_C \leq l \leq k_C} \sum_{i+j=l, h_A \leq i \leq k_A, h_B \leq j \leq k_B} a_i b_j \varepsilon^l \\
&+ \sum_{k_C < i+j, h_A \leq i \leq k_A, h_B \leq j \leq k_B} a_i b_j \varepsilon^{i+j} + \sum_{h_B \leq j \leq k_B} b_j \varepsilon^j o_A(\varepsilon^{k_A}) \\
&+ \sum_{h_A \leq i \leq k_A} a_i \varepsilon^i o_B(\varepsilon^{k_B}) + o_A(\varepsilon^{k_A}) o_B(\varepsilon^{k_B}) \\
&= \sum_{h_C \leq l \leq k_C} c_l \varepsilon^l + o_C(\varepsilon^{k_C}), \tag{8}
\end{aligned}$$

where

$$\begin{aligned}
o_C(\varepsilon^{k_C}) &= \sum_{k_C < i+j, h_A \leq i \leq k_A, h_B \leq j \leq k_B} a_i b_j \varepsilon^{i+j} + \sum_{h_B \leq j \leq k_B} b_j \varepsilon^j o_A(\varepsilon^{k_A}) \\
&+ \sum_{h_A \leq i \leq k_A} a_i \varepsilon^i o_B(\varepsilon^{k_B}) + o_A(\varepsilon^{k_A}) o_B(\varepsilon^{k_B}). \tag{9}
\end{aligned}$$

Obviously, $\frac{o_C(\varepsilon^{k_C})}{\varepsilon^{k_C}} \rightarrow 0$ as $\varepsilon \rightarrow 0$. It should be noted that the accumulation of coefficients for powers ε^l can be made in (8) only up to the maximal value $l = k_C = (k_A + h_B) \wedge (k_B + h_A)$, because of the presence in the expression for remainder $o_C(\varepsilon^{k_C})$ terms $b_{h_B} \varepsilon^{h_B} o_A(\varepsilon^{k_A})$ and $a_{h_A} \varepsilon^{h_A} o_B(\varepsilon^{k_B})$.

Also, relation (9) readily implies relations **(a)** – **(c)**, which determines parameters $\delta_C, G_C, \varepsilon_C$ in proposition **(iii)** of Lemma 2.

The assumptions of proposition **(iv)** in Lemma 2* imply that $\varepsilon^{-h_B} B(\varepsilon) \rightarrow b_{h_B} \neq 0$ as $\varepsilon \rightarrow 0$. This relation implies that there exists $0 < \varepsilon'_0 \leq \varepsilon_0$ such that $B(\varepsilon) \neq 0$ for $\varepsilon \in (0, \varepsilon'_0]$, and, thus, function $C(\varepsilon) = \frac{1}{B(\varepsilon)}$ is well defined for $\varepsilon \in (0, \varepsilon'_0]$.

Note that $h_B \leq k_B$. The assumptions of proposition **(iv)** of Lemma 2* imply that, $\varepsilon^{h_B} C(\varepsilon) = (b_{h_B} + \dots + b_{k_B} \varepsilon^{k_B - h_B} + o_B(\varepsilon^{h_B}) \varepsilon^{-h_B})^{-1} \rightarrow b_{h_B}^{-1} = c_{h_C}$ as $\varepsilon \rightarrow 0$. This relation means that function $\varepsilon^{h_B} C(\varepsilon)$ can be represented in the form $\varepsilon^{h_B} C(\varepsilon) = c_{h_C} + o(1)$, where $c_{h_C} = b_{h_B}^{-1}$, or, equivalently, that the following representation takes place, $C(\varepsilon) = c_{h_C} \varepsilon^{-h_B} + o_1(\varepsilon^{-h_B})$, $\varepsilon \in (0, \varepsilon'_0]$, where $\frac{o_1(\varepsilon^{-h_B})}{\varepsilon^{-h_B}} \rightarrow 0$ as $\varepsilon \rightarrow 0$.

The latter two relations prove proposition **(iv)** of Lemma 2*, for the case $h_B = k_B$. Indeed, these relations mean that function $C(\varepsilon) = \frac{1}{B(\varepsilon)}$ can be represented in the form of (h_C, k_C) -expansion with parameters $h_C = -h_B$, $k_C = k_B - 2h_B = -h_B = h_C$ and coefficient $c_{h_C} = b_{h_B}^{-1}$. Moreover, since $B(\varepsilon) \cdot C(\varepsilon) \equiv 1$, $0 < \varepsilon \leq \varepsilon'_0$, remainder $c_1(\varepsilon)$ can be found from the following

relation, $(b_{h_B}\varepsilon^{h_B} + o(\varepsilon^{h_B}))(c_{h_C}\varepsilon^{-h_B} + o_1(\varepsilon^{-h_B})) \equiv 1$ that yields formula, $o_1(\varepsilon^{-h_B}) = -\frac{c_{h_C}\varepsilon^{-h_B}o_B(\varepsilon^{h_B})}{b_{h_B}\varepsilon^{h_B}+o_B(\varepsilon^{h_B})}$. This is formula (c) from proposition (iv) of Lemma 2*, for the case $h_B = k_B$. Note that, in the case $h_B = k_B$, the above asymptotic expansion for function $C(\varepsilon)$ can not be extended. Indeed, $\varepsilon^{h_B-1}o_1(\varepsilon^{-h_B}) = \varepsilon^{h_B-1}(C(\varepsilon) - c_{h_C}\varepsilon^{-h_B}) = -\frac{c_{h_C}}{b_{h_B}+o_B(\varepsilon^{h_B})\varepsilon^{-h_B}}\frac{o_B(\varepsilon^{h_B})\varepsilon^{-h_B}}{\varepsilon}$. The term $\frac{o_B(\varepsilon^{h_B})\varepsilon^{-h_B}}{\varepsilon}$ on the right hand side in the latter relation has an uncertain asymptotic behavior as $\varepsilon \rightarrow 0$.

Let us now assume that $h_B + 1 \leq k_B$. In this case, the assumptions of proposition (iv) of Lemma 2* and the above asymptotic relations imply that $\varepsilon^{h_B-1}o_1(\varepsilon^{-h_B}) = \varepsilon^{h_B-1}(C(\varepsilon) - c_{h_C}\varepsilon^{-h_B}) = (b_{h_B} + \dots + b_{k_B}\varepsilon^{k_B-h_B} + o_B(\varepsilon^{k_B})\varepsilon^{-h_B})^{-1}(-b_{h_B+1}c_{h_C} - \dots - b_{k_B}c_{h_C}\varepsilon^{k_B-h_B-1} - o_B(\varepsilon^{k_B})c_{h_C}\varepsilon^{-h_B-1}) \rightarrow -b_{h_B}^{-1}b_{h_B+1}c_{h_C} = c_{h_C+1}$ as $\varepsilon \rightarrow 0$. This relation means that function $\varepsilon^{h_B-1} \cdot o_1(\varepsilon^{-h_B})$ can be represented in the form $\varepsilon^{h_B-1}o_1(\varepsilon^{-h_B}) = c_{h_C+1} + o(1)$, where $c_{h_C+1} = -b_{h_B}^{-1}b_{h_B+1}c_{h_C}$, or, equivalently, that the following representation takes place, $C(\varepsilon) = c_{h_C}\varepsilon^{-h_B} + c_{h_C+1}\varepsilon^{-h_B+1} + o_2(\varepsilon^{-h_B+1})$, $\varepsilon \in (0, \varepsilon'_0]$, where $\frac{o_2(\varepsilon^{-h_B+1})}{\varepsilon^{-h_B+1}} \rightarrow 0$ as $\varepsilon \rightarrow 0$.

The latter two relations prove proposition (iv) of Lemmas 2*, for the case $h_B + 1 = k_B$. Indeed, these relations mean that function $C(\varepsilon)$ can be represented in the form of (h_C, k_C) -expansion with parameters $h_C = -h_B$, $k_C = k_B - 2h_B = -h_B + 1 = h_C + 1$ and coefficients $c_{h_C} = b_{h_B}^{-1}$, $c_{h_C+1} = -b_{h_B}^{-1}b_{h_B+1}c_{h_C}$. Moreover, since $B(\varepsilon) \cdot C(\varepsilon) \equiv 1$, the remainder $o_2(\varepsilon^{-h_B+1})$ can be found from the following relation, $(b_{h_B}\varepsilon^{h_B} + b_{h_B+1}\varepsilon^{h_B+1} + o(\varepsilon^{h_B+1})) \cdot (c_{h_C}\varepsilon^{-h_B} + c_{h_C+1}\varepsilon^{-h_B+1} + o_2(\varepsilon^{-h_B+1})) \equiv 1$ that yields formula, $o_2(\varepsilon^{-h_B+1}) = -\frac{b_{h_B+1}c_{h_C+1}\varepsilon^2 + (c_{h_C}\varepsilon^{-h_B} + c_{h_C+1}\varepsilon^{-h_B+1})o_B(\varepsilon^{h_B+1})}{b_{h_B}\varepsilon^{h_B} + o_B(\varepsilon^{h_B})}$. This is formula (c) from proposition (iv) of Lemma 2*, for the case $h_B + 1 = k_B$. Note that, in the case $h_B + 1 = k_B$, the above asymptotic expansion for function $C(\varepsilon)$ can not be extended. Indeed, $\varepsilon^{h_B-2}C_2(\varepsilon) = \varepsilon^{h_B-2}(C(\varepsilon) - c_{h_C}\varepsilon^{-h_B} - c_{h_C+1}\varepsilon^{-h_B+1}) = -\frac{c_{h_C}}{b_{h_B}+b_{h_B+1}\varepsilon+o_B(\varepsilon^{h_B+1})\varepsilon^{-h_B}} \times \frac{o_B(\varepsilon^{h_B+1})\varepsilon^{-h_B-1}}{\varepsilon}$. The term $\frac{o_B(\varepsilon^{h_B+1})\varepsilon^{-h_B-1}}{\varepsilon}$ on the right hand side in the latter relation has an uncertain asymptotic behavior as $\varepsilon \rightarrow 0$.

We can repeat the above arguments for the general case $h_B + n = k_B$, for any $n = 0, 1, \dots$ and to prove that, in the case $h_B + n = k_B$, function $C(\varepsilon)$ can be represented in the form of (h_C, k_C) -expansion with parameters $h_C = -h_B$, $k_C = k_B - 2h_B = -h_B + n = h_C + n$ and coefficients $c_{h_C}, \dots, c_{k_C}$ given in proposition (iv) of Lemma 2*. Moreover, identity $B(\varepsilon) \cdot C(\varepsilon) \equiv$

$1, 0 < \varepsilon \leq \varepsilon'_0$, let us find the corresponding remainder $o_C(\varepsilon^{k_C})$ from the following relation,

$$(b_{h_B}\varepsilon^{h_B} + \dots + b_{k_B}\varepsilon^{k_B} + o_B(\varepsilon^{k_B})) \times (c_{h_C}\varepsilon^{h_C} + \dots + c_{k_C}\varepsilon^{k_C} + o_C(\varepsilon^{k_C})) \equiv 1. \quad (10)$$

Proposition **(iii)** of Lemma 2_{*}, applied to the product on the left hand side in relation (10), permits to represent this product in the form of (h, k) -expansion with parameters $h = h_B + h_C = h_B - h_B = 0$ and $k = (k_B + h_C) \wedge (h_B + k_C) = (k_B - h_B) \wedge (k_B - 2h_B + h_B) = k_B - h_B$. By canceling coefficient for ε^l on the left and right hand sides in relation (10), for $l = 0, \dots, k_B - h_B$, and then, by solving equation (10) with respect to the remainder $o_C(\varepsilon^{k_C})$, we get the formula for this remainder given in proposition **(iv)** of Lemma 2_{*},

$$\begin{aligned} o_C(\varepsilon^{k_C}) &= - \frac{\sum_{k_B - h_B < i+j, h_B \leq i \leq k_B, h_C \leq j \leq k_C} b_i c_j \varepsilon^{i+j} + \sum_{h_C \leq j \leq k_C} c_j \varepsilon^j o_B(\varepsilon^{k_B})}{b_{h_B}\varepsilon^{h_B} + \dots + b_{k_B}\varepsilon^{k_B} + o_B(\varepsilon^{k_B})} \\ &= - \frac{\sum_{k_B - h_B < i+j, h_B \leq i \leq k_B, h_C \leq j \leq k_C} b_i c_j \varepsilon^{i+j-h_B}}{b_{h_B} + \dots + b_{k_B}\varepsilon^{k_B-h_B} + o_B(\varepsilon^{k_B})\varepsilon^{h_B}} \\ &\quad - \frac{\sum_{h_C \leq j \leq k_C} c_j \varepsilon^{j-h_B} o_B(\varepsilon^{k_B})}{b_{h_B} + \dots + b_{k_B}\varepsilon^{k_B-h_B} + o_B(\varepsilon^{k_B})\varepsilon^{-h_B}}. \end{aligned} \quad (11)$$

The assumptions made in proposition **(iv)** of Lemma 2, imply that $B(\varepsilon) \neq 0$ and the following inequality holds for $0 < \varepsilon \leq \varepsilon_C$, where ε_C is given in proposition **(iv)** of Lemma 2,

$$\begin{aligned} &|b_{h_B} + b_{h_B+1}\varepsilon + \dots + b_{k_B}\varepsilon^{k_B-h_B} + o_B(\varepsilon^{k_B})\varepsilon^{-h_B}| \\ &\geq |b_{h_B}| - (|b_{h_B+1}|\varepsilon + \dots + |b_{k_B}|\varepsilon^{k_B-h_B} + G_B\varepsilon^{k_B-h_B+\delta_B}) \\ &\geq \frac{|b_{h_B}|}{2} > 0. \end{aligned} \quad (12)$$

The existence of ε'_0 declared in proposition **(iv)** of Lemma 2 is obvious. For example, one can choose $\varepsilon'_0 = \varepsilon_C$. It is also useful to note that formulas given in proposition **(iv)** of Lemma 2 imply that $\varepsilon_C = \varepsilon_B \wedge \tilde{\varepsilon}_B \in (0, \varepsilon_0]$, since $\varepsilon_B \in (0, \varepsilon_0]$ and $\tilde{\varepsilon}_B \in (0, \infty)$.

The assumptions made in proposition **(iv)** of Lemma 2 and inequality

(12) imply that the following inequality holds, for $0 < \varepsilon \leq \varepsilon_C$,

$$\begin{aligned}
|o_C(\varepsilon^{k_C})| &\leq \varepsilon^{k_B-2h_B+\delta_B} \left(\frac{|b_{h_B}|}{2}\right)^{-1} \\
&\times \left(\sum_{k_B-h_B < i+j, h_B \leq i \leq k_B, h_C \leq j \leq k_C} |b_i| |c_j| \varepsilon_C^{i+j-k_B+h_B-\delta_B} \right. \\
&\left. + G_B \sum_{h_C \leq j \leq k_C} |c_j| \varepsilon_C^{j+h_B} \right). \tag{13}
\end{aligned}$$

Inequality (13) proofs proposition **(iv)** of Lemma 2.

Propositions **(v)** of Lemmas 2_{*} and 2 and relations **(a)** – **(c)** given in these propositions can be obtained by direct application, respectively, of propositions **(iii)** and **(iv)** of Lemmas 2_{*} and 2, to the product $D(\varepsilon) = A(\varepsilon) \cdot \frac{1}{B(\varepsilon)}$.

Now, when it is already known that $D(\varepsilon) = A(\varepsilon) \cdot \frac{1}{B(\varepsilon)}$ is a (h_D, k_D) -expansion, with parameters $h_D = h_A - h_B$ and $k_D = (k_A - h_B) \wedge (k_B - 2h_B + h_A)$, multiplication of $D(\varepsilon)$ by $B(\varepsilon)$ yields the following relation holding for $\varepsilon \in (0, \varepsilon'_0]$,

$$\begin{aligned}
A(\varepsilon) &= D(\varepsilon)B(\varepsilon) = a_{h_A}\varepsilon^{h_A} + \dots + a_{h_A}\varepsilon^{k_A} + o_A(\varepsilon^{k_A}) \\
&= (d_{h_D}\varepsilon^{h_D} + \dots + d_{h_D}\varepsilon^{k_D} + o_D(\varepsilon^{k_D})) \\
&\quad \times (b_{h_B}\varepsilon^{h_B} + \dots + b_{h_B}\varepsilon^{k_B} + o_B(\varepsilon^{k_B})). \tag{14}
\end{aligned}$$

By equating coefficients for powers ε^l for $l = h_D, \dots, k_D$ on the left and right hand sides of the third equality in relation (14), we get alternative formulas **(e)** for coefficients $d_{h_A}, \dots, d_{k_D}$ given in proposition **(v)** of Lemma 2_{*}.

Proposition **(iii)** of Lemma 2, applied to the product on the right hand side in (14), permits to represent this product in the form of (h, k) -expansion with parameters $h = h_B + h_D = h_B + h_A - h_B = h_A$ and $k = (k_D + h_B) \wedge (k_B + h_D) = ((k_A - h_B) \wedge (k_B - 2h_B + h_A) + h_B) \wedge (k_B + h_A - h_B) = k_A \wedge (k_B + h_A - h_B)$. By canceling coefficient for ε^l on the left and right hand sides in relation (14), for $l = h_A, \dots, k_A \wedge (k_B + h_A - h_B)$, and then, by solving equation (14) with respect to the remainder $o_D(\varepsilon^{k_D})$, we get the formula **(f)** for this remainder given in proposition **(v)** of Lemma 2_{*},

$$o_D(\varepsilon^{k_D}) = \frac{\sum_{k_A \wedge (k_B + h_A - h_B) < l \leq k_A} a_l \varepsilon^l + o_A(\varepsilon^{k_A})}{b_{h_B}\varepsilon^{h_B} + \dots + b_{k_B}\varepsilon^{k_B} + o_B(\varepsilon^{k_B})}$$

$$\begin{aligned}
& - \frac{\sum_{k_A \wedge (k_B + h_A - h_B) < i+j, h_B \leq i \leq k_B, h_D \leq j \leq k_D} b_i d_j \varepsilon^{i+j}}{b_{h_B} \varepsilon^{h_B} + \dots + b_{k_B} \varepsilon^{k_B} + o_B(\varepsilon^{k_B})} \\
& - \frac{\sum_{h_D \leq j \leq k_D} d_j \varepsilon^j o_B(\varepsilon^{k_B})}{b_{h_B} \varepsilon^{h_B} + \dots + b_{k_B} \varepsilon^{k_B} + o_B(\varepsilon^{k_B})} \\
& = \frac{\sum_{k_A \wedge (k_B + h_A - h_B) < l \leq k_A} a_l \varepsilon^{l-h_B} + o_A(\varepsilon^{k_A}) \varepsilon^{-h_B}}{b_{h_B} + \dots + b_{k_B} \varepsilon^{k_B-h_B} + o_B(\varepsilon^{k_B}) \varepsilon^{-h_B}} \\
& - \frac{\sum_{k_A \wedge (k_B + h_A - h_B) < i+j, h_B \leq i \leq k_B, h_D \leq j \leq k_D} b_i d_j \varepsilon^{i+j-h_B}}{b_{h_B} + \dots + b_{k_B} \varepsilon^{k_B-h_B} + o_B(\varepsilon^{k_B}) \varepsilon^{-h_B}} \\
& - \frac{\sum_{h_D \leq j \leq k_D} d_j \varepsilon^{j-h_B} o_B(\varepsilon^{k_B})}{b_{h_B} + \dots + b_{k_B} \varepsilon^{k_B-h_B} + o_B(\varepsilon^{k_B}) \varepsilon^{-h_B}}. \tag{15}
\end{aligned}$$

Inequality (12) and the assumptions made in proposition **(v)** of Lemma 2 finally imply that the following inequality holds, for $0 < \varepsilon \leq \varepsilon_D$ given in relation **(f)** of this proposition,

$$\begin{aligned}
|o_D(\varepsilon^{k_D})| & \leq \varepsilon^{k_D+\delta_D} \left(\frac{|b_{h_B}|}{2} \right)^{-1} \left(\sum_{k_A \wedge (k_B + h_A - h_B) < l \leq k_A} |a_l| \varepsilon_D^{l-k_D-h_B-\delta_D} \right. \\
& + \sum_{k_A \wedge (k_B + h_A - h_B) < i+j, h_B \leq i \leq k_B, h_D \leq j \leq k_D} |b_i| |d_j| \varepsilon_D^{i+j-k_D-h_B-\delta_D} \\
& + G_A e_D^{k_A+\delta_A-h_B-k_D-\delta_D} \\
& \left. + G_B \sum_{h_D \leq j \leq k_D} |d_j| \varepsilon_D^{j+k_B+\delta_B-h_B-k_D-\delta_D} \right). \tag{16}
\end{aligned}$$

Inequality (16) yields relations **(d)** – **(f)** given in proposition **(v)** of Lemma 2.

A.3. Lemma 3_{*} is a direct corollary of Lemma 2_{*}. Proofs of propositions **(i)** and **(ii)** in Lemma 3 are analogous to proofs of the corresponding propositions in Lemma 2. Proposition **(iii)** of Lemma 3 is obvious.

A.4. The first two identities for Laurent asymptotic expansions given in proposition **(i)** of Lemma 4_{*} are obvious. The third identity given in this proposition follows in an obvious way from proposition **(i)** of Lemma 2_{*}. By applying propositions **(iii)** and **(iv)** of Lemma 2_{*} to the product $C(\varepsilon) = A(\varepsilon) \cdot A(\varepsilon)^{-1}$, we get parameters $h_C = h_{A \cdot A^{-1}} = h_A - h_A = 0$, $k_C = k_{A \cdot A^{-1}} = (k_A - h_A) \wedge (k_A - 2h_A + h_A) = k_A - h_A$ and coefficients $c_n = I(n = 0)$, $n = 0, \dots, k_C$. Also, relations (10) and (11) imply that the elimination identity $A(\varepsilon) \cdot A(\varepsilon)^{-1} \equiv 1$ holds, since the remainder of Laurent asymptotic

expansion for function $A(\varepsilon)^{-1}$ is given by formula (c) from proposition (iv) of Lemma 2*.

Propositions (ii) and (iii) of Lemma 4* in the parts concerned commutative property of summation and multiplication operations follow from, respectively, propositions (ii) and (iii) of Lemma 2*.

Let $D(\varepsilon) = (A(\varepsilon) + B(\varepsilon)) + C(\varepsilon) = A(\varepsilon) + (B(\varepsilon) + C(\varepsilon))$. Using propositions (ii) of Lemma 2*, we get, $h_D = h_{(A+B)+C} = (h_A \wedge h_B) \wedge h_C = h_A \wedge (h_B \wedge h_C) = h_{A+(B+C)}$ and $k_D = k_{(A+B)+C} = (k_A \wedge k_B) \wedge k_C = k_A \wedge (k_B \wedge k_C) = k_{A+(B+C)}$. These relations and Lemma 1* imply equalities for the corresponding coefficients and remainders, for the asymptotic expansions of functions $(A(\varepsilon) + B(\varepsilon)) + C(\varepsilon)$ and $A(\varepsilon) + (B(\varepsilon) + C(\varepsilon))$. The above remarks prove proposition (ii) of Lemma 4* in the part concerned with the associative property of summation operation for Laurent asymptotic expansions.

Let $D(\varepsilon) = (A(\varepsilon) \cdot B(\varepsilon)) \cdot C(\varepsilon) = A(\varepsilon) \cdot (B(\varepsilon) \cdot C(\varepsilon))$. Using propositions (iii) of Lemma 2*, we get, $h_D = h_{(A \cdot B) \cdot C} = h_{A \cdot B} + h_C = h_A + h_B + h_C = h_A + h_{B \cdot C} = h_{A \cdot (B \cdot C)}$ and $k_D = k_{(A \cdot B) \cdot C} = (k_{A \cdot B} + h_C) \wedge (k_C + h_{A \cdot B}) = ((k_A + h_B) \wedge (k_B + h_A)) \wedge h_C \wedge (k_C + (h_A + h_B)) = (k_A + h_B + h_C) \wedge (k_B + h_A + h_C) \wedge (k_C + h_A + h_B) = (k_A + (h_B + h_C)) \wedge ((k_B + h_C) \wedge (k_C + h_B)) + h_A = (k_A + h_{B \cdot C}) \wedge (k_{B \cdot C} + h_A) = k_{A \cdot (B \cdot C)}$. These relations and Lemma 1* imply equalities for the corresponding coefficients and remainders, for the asymptotic expansions of functions $(A(\varepsilon) \cdot B(\varepsilon)) \cdot C(\varepsilon)$ and $A(\varepsilon) \cdot (B(\varepsilon) \cdot C(\varepsilon))$. The above remarks prove proposition (iii) of Lemma 4* in the part concerned with the associative property of multiplication operation for Laurent asymptotic expansions.

Let $D(\varepsilon) = (A(\varepsilon) + B(\varepsilon)) \cdot C(\varepsilon) = A(\varepsilon) \cdot C(\varepsilon) + B(\varepsilon) \cdot C(\varepsilon)$. Using propositions (ii) and (iii) of Lemma 2*, we get, $h_D = h_{(A+B) \cdot C} = h_{A+B} + h_C = h_A \wedge h_B + h_C = (h_A + h_C) \wedge (h_B + h_C) = h_{A \cdot C} \wedge h_{B \cdot C} = h_{A \cdot C + B \cdot C}$ and $k_D = k_{(A+B) \cdot C} = (k_{A+B} + h_C) \wedge (k_C + h_{A+B}) = (k_A \wedge k_B + h_C) \wedge (k_C + h_A \wedge h_B) = (k_A + h_C) \wedge (k_B + h_C) \wedge (k_C + h_A) \wedge (k_C + h_B) = ((k_A + h_C) \wedge (k_C + h_A)) \wedge ((k_B + h_C) \wedge (k_C + h_B)) = k_{A \cdot C} \wedge k_{B \cdot C} = k_{A \cdot C + B \cdot C}$. These relations and Lemma 1* imply equalities for the corresponding coefficients and remainders, for the asymptotic expansions of functions $(A(\varepsilon) + B(\varepsilon)) \cdot C(\varepsilon)$ and $A(\varepsilon) \cdot C(\varepsilon) + B(\varepsilon) \cdot C(\varepsilon)$. The above remarks prove proposition (iv) of Lemma 4* concerned with the distributive property of summation and multiplication operations for Laurent asymptotic expansions.

Propositions (i) and (ii) of Lemma 4 readily follow from, respectively, propositions (ii) and (iii) of Lemma 2. Finally, proposition (iii) of Lemma 4 follows from relations $\delta_{cA} = \delta_A$, δ_{A+B} , $\delta_{A \cdot B}$, $\delta_{A/B} \geq \delta_A \wedge \delta_B$ and $\delta_{A_1 + \dots + A_N}$,

$\delta_{A_1 \times \dots \times A_N} \geq \min_{1 \leq m \leq N} \delta_{A_m}$, given, respectively, in Lemmas 2 and 3. $\square$

Appendix B: Examples

Let us, first, comment some general questions connected with construction of examples illustrating the asymptotic results presented in the paper.

Let $\mathbb{Y}_i \neq \emptyset, i \in \mathbb{X}$ be some subsets of space $\mathbb{X}$ such that condition **A (c)** holds for these sets, i.e., for every pair of states $i, j \in \mathbb{X}$, there exists an integer $n_{ij} \geq 1$ and a chain of states $i = l_{ij,0}, l_{ij,1}, \dots, l_{ij,n_{ij}} = j$ such that $l_{ij,1} \in \mathbb{Y}_{l_{ij,0}}, \dots, l_{ij,n_{ij}} \in \mathbb{Y}_{l_{ij,n_{ij}-1}}$.

Let us also choose some $\varepsilon_0 \in (0, 1]$.

We define $p_{ij}(\varepsilon) = 0, \varepsilon \in (0, \varepsilon_0], j \in \overline{\mathbb{Y}}_i, i \in \mathbb{X}$, i.e., assume that condition **A (b)** holds.

Let $p_{ij}(\varepsilon), \varepsilon \in (0, \varepsilon_0], j \in \mathbb{Y}_i, i \in \mathbb{X}$ be some real-valued functions which satisfy condition **D**, i.e., can be represented in the form of Taylor asymptotic expansions, $p_{ij}(\varepsilon) = \sum_{l=l_{ij}^-}^{l_{ij}^+} a_{ij}[l] \varepsilon^l + o_{ij}(\varepsilon^{l_{ij}^+}), \varepsilon \in (0, \varepsilon_0]$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$, where **(a)** $a_{ij}[l_{ij}^-] > 0$ and $0 \leq l_{ij}^- \leq l_{ij}^+ < \infty$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$; **(b)** $o_{ij}(\varepsilon^{l_{ij}^+})/\varepsilon^{l_{ij}^+} \rightarrow 0$ as $\varepsilon \rightarrow 0$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$.

Condition **D** does not guarantee that matrix $\|p_{ij}(\varepsilon)\|$ is stochastic, for every $\varepsilon \in (0, \varepsilon_0]$. This can be achieved by imposing some additional conditions on coefficients and remainders in the above asymptotic expansions.

Let us choose arbitrary numbers $0 < \alpha_{ij} < \frac{1}{2}$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$.

Condition **D (b)** guarantees that, for every $j \in \mathbb{Y}_i, i \in \mathbb{X}$, there exists $\varepsilon_{\alpha_{ij}, ij} \in (0, \varepsilon_0]$ such that, for $\varepsilon \in (0, \varepsilon_{\alpha_{ij}, ij}]$,

$$|o_{ij}(\varepsilon^{l_{ij}^+})/\varepsilon^{l_{ij}^+}| \leq \alpha_{ij} |a_{ij}[l_{ij}^-]|. \quad (17)$$

It is useful to note that in the case, where condition **D'** holds, an explicit value for parameters $\varepsilon_{\alpha_{ij}, ij}$ can be derived from the inequalities, $|o_{ij}(\varepsilon^{l_{ij}^+})| \leq G_{ij} \varepsilon^{l_{ij}^+ + \delta_{ij}}, \varepsilon \in (0, \varepsilon_{ij}], j \in \mathbb{Y}_i, i \in \mathbb{X}$, penetrating this condition. These inequalities imply that relation (17) holds for $\varepsilon_{\alpha_{ij}, ij} = \varepsilon_{ij} \wedge \left(\frac{\alpha_{ij} |a_{ij}[l_{ij}^-]|}{G_{ij}}\right)^{\frac{1}{\delta_{ij}}}$, $j \in \mathbb{Y}_i, i \in \mathbb{X}$.

Let us also define $A_{\varepsilon_0, ij} = \sum_{l_{ij}^- < l \leq l_{ij}^+} |a_{ij}[l]| \varepsilon_0^{l - l_{ij}^- - 1}, j \in \mathbb{Y}_i, i \in \mathbb{X}$ and $\varepsilon'_{\alpha_{ij}, ij} = \varepsilon_{\alpha_{ij}, ij}$ if $A_{\varepsilon_0, ij} = 0$ or $\varepsilon'_{\alpha_{ij}, ij} = \varepsilon_{\alpha_{ij}, ij} \wedge \frac{\alpha_{ij} |a_{ij}[l_{ij}^-]|}{A_{\varepsilon_0, ij}}$ if $A_{\varepsilon_0, ij} > 0$.

The following inequality holds, for every $\varepsilon \in (0, \varepsilon'_{\alpha_{ij}, ij}]$ and $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$,

$$\begin{aligned} p_{ij}(\varepsilon) &\geq \varepsilon^{l_{ij}^-} (a_{ij}[l_{ij}^-] - \varepsilon A_{\varepsilon_0, ij} - \varepsilon_0^{l_{ij}^+ - l_{ij}^-} |o_{ij}(\varepsilon^{l_{ij}^+}) / \varepsilon^{l_{ij}^+}|) \\ &\geq \varepsilon^{l_{ij}^-} a_{ij}[l_{ij}^-] (1 - 2\alpha_{ij}) > 0. \end{aligned} \quad (18)$$

Let us now define $\varepsilon'_0 = \min_{j \in \mathbb{Y}_i, i \in \mathbb{X}} \varepsilon'_{\alpha_{ij}, ij}$.

Obviously, $p_{ij}(\varepsilon) > 0$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$, $\varepsilon \in (0, \varepsilon'_0]$. Thus, conditions **A (a) – (b)** hold, if parameter ε_0 is replaced by the new value ε'_0 .

The question about holding of the stochasticity relation $\sum_{j \in \mathbb{Y}_i} p_{ij}(\varepsilon) = 1$, $\varepsilon \in (0, \varepsilon_0]$, $i \in \mathbb{X}$ is more complex. According Lemma 5*, under conditions **A (a) – (b)** and **D**, the above stochasticity relation is equivalent to condition **F** formulated in Section 3*.

First, condition **F** requires holding of the following relation,

$$\sum_{j \in \mathbb{Y}_i} a_{ij}[l] = \mathbf{I}(l = 0), \quad 0 \leq l \leq l_{i, \mathbb{Y}_i}^+, \quad i \in \mathbb{X}, \quad (19)$$

where (a) $l_{i, \mathbb{Y}_i}^\pm = \min_{j \in \mathbb{Y}_i} l_{ij}^\pm$, $i \in \mathbb{X}$ and (b) $a_{ij}[l] = 0$, for $0 \leq l < l_{ij}^-, j \in \mathbb{Y}_i$, $i \in \mathbb{X}$.

Note that relation (19) implies that parameters $l_{i, \mathbb{Y}_i}^- = 0$, $i \in \mathbb{X}$.

It is not difficult to choose coefficients $a_{ij}[l]$, $l = l_{ij}^- \leq l \leq l_{ij}^+$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$ in such way that relation (19) would hold. Any such coefficients, with the first coefficients $a_{ij}[l_{ij}^-] > 0$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$, can serve as coefficients in the asymptotic expansions penetrating condition **D**.

Second, condition **F** requires holding of the following identity, for every $i \in \mathbb{X}$,

$$\sum_{j \in \mathbb{Y}_i} \left(\sum_{l_{i, \mathbb{Y}_i}^+ < l \leq l_{ij}^+} a_{ij}[l] \varepsilon^l + o_{ij}(\varepsilon^{l_{ij}^+}) \right) \equiv 0. \quad (20)$$

Remainders $o_{ij}(\varepsilon^{l_{ij}^+})$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$ satisfying the above identities can be chosen in different ways.

The simplest one is to choose $o_{ij}(\varepsilon^{l_{ij}^+}) \equiv 0$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$. In this case, the above identities would reduce to equalities, $\sum_{j \in \mathbb{Y}_i} a_{ij}[l] = 0$, $l_{i, \mathbb{Y}_i}^+ < l \leq l_{i, \mathbb{Y}_i}^*$, $i \in \mathbb{X}$, where (a) $l_{i, \mathbb{Y}_i}^* = \max_{j \in \mathbb{Y}_i} l_{ij}^+$, $i \in \mathbb{X}$ and (b) $a_{ij}[l] = 0$, for $l_{ij}^+ < l \leq l_{i, \mathbb{Y}_i}^*$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$. These equalities supplement equalities given in relation (19). Such choice of remainders corresponds to models with polynomial perturbations.

We, however, would like to impose on remainders conditions mainly required of them by conditions **D** or **D'**.

There always exist $j_i \in \mathbb{Y}_i, i \in \mathbb{X}$ such that $l_{ij_i}^+ = l_{i, \mathbb{Y}_i}^+, i \in \mathbb{X}$.

Identity (20) can be rewritten in the following form, for every $i \in \mathbb{X}$,

$$o_{ij_i}(\varepsilon^{l_{ij_i}^+}) \equiv - \sum_{j \in \mathbb{Y}_i, j \neq j_i} \left(\sum_{l_{i, \mathbb{Y}_i}^+ < l \leq l_{ij}^+} a_{ij}[l] \varepsilon^l + o_{ij}(\varepsilon^{l_{ij}^+}) \right). \quad (21)$$

Relation (21) can be used as the formula defining remainders $o_{ij_i}(\varepsilon^{l_{ij_i}^+}), i \in \mathbb{X}$, via remainders $o_{ij}(\varepsilon^{l_{ij}^+}), j \in \mathbb{Y}_i, j \neq j_i, i \in \mathbb{X}$ penetrating the corresponding asymptotic expansions in condition **D**.

Since $l_{ij_i}^+ = l_{i, \mathbb{Y}_i}^+, i \in \mathbb{X}$, the following relation holds, for remainder $o_{ij}(\varepsilon^{l_{ij}^+})$ defined by relation (21), for every $i \in \mathbb{X}$,

$$o_{ij_i}(\varepsilon^{l_{ij_i}^+}) / \varepsilon^{l_{ij_i}^+} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \quad (22)$$

Thus, remainders $o_{ij_i}(\varepsilon^{l_{ij_i}^+}), i \in \mathbb{X}$ defined by relation (21) can also serve in the corresponding asymptotic expansions in condition **D**.

Moreover, let us assume that remainders $o_{ij}(\varepsilon^{l_{ij}^+}), j \in \mathbb{Y}_i, j \neq j_i, i \in \mathbb{X}$ satisfy the inequalities, $|o_{ij}(\varepsilon^{l_{ij}^+})| \leq G_{ij} \varepsilon^{l_{ij}^+ + \delta_{ij}}, \varepsilon \in (0, \varepsilon_{ij}], j \in \mathbb{Y}_i, j \neq j_i, i \in \mathbb{X}$, penetrating condition **D'**.

Let us define $\varepsilon_{ij_i} = \min_{j \in \mathbb{Y}_i, j \neq j_i} \varepsilon_{ij}, i \in \mathbb{X}$ and $\delta_{ij_i} = \min_{j \in \mathbb{Y}_i, j \neq j_i} \delta_{ij}, i \in \mathbb{X}$.

In this case, the following inequality holds, for every $\varepsilon \in (0, \varepsilon_{ij_i}], i \in \mathbb{X}$,

$$\begin{aligned} |o_{ij_i}(\varepsilon^{l_{ij_i}^+})| &\leq \sum_{j \in \mathbb{Y}_i, j \neq j_i} \left(\sum_{l_{i, \mathbb{Y}_i}^+ < l \leq l_{ij}^+} |a_{ij}[l]| \varepsilon^l + |o_{ij}(\varepsilon^{l_{ij}^+})| \right) \\ &\leq \left(\sum_{j \in \mathbb{Y}_i, j \neq j_i} \left(\sum_{l_{i, \mathbb{Y}_i}^+ < l \leq l_{ij}^+} |a_{ij}[l]| \varepsilon_0^{l - l_{ij_i}^+ - \delta_{ij_i}} \right. \right. \\ &\quad \left. \left. + \varepsilon_0^{l_{ij}^+ - l_{ij_i}^+ + \delta_{ij} - \delta_{ij_i}} G_{ij} \right) \varepsilon^{l_{ij_i}^+ + \delta_{ij_i}} \right) = G_{ij_i} \varepsilon^{l_{ij_i}^+ + \delta_{ij_i}}. \end{aligned} \quad (23)$$

Thus, the inequalities, $|o_{ij_i}(\varepsilon^{l_{ij_i}^+})| \leq G_{ij_i} \varepsilon^{l_{ij_i}^+ + \delta_{ij_i}}, \varepsilon \in (0, \varepsilon_{ij_i}], i \in \mathbb{X}$, penetrating condition **D'** hold for remainders $o_{ij_i}(\varepsilon^{l_{ij_i}^+}), i \in \mathbb{X}$, with parameters $\varepsilon_{ij_i}, \delta_{ij_i}$ and G_{ij_i} defined above.

As follows from the above remarks, identity (21) holds for remainders $o_{ij}(\varepsilon^{l_{ij}^+}), j \in \mathbb{Y}_i, i \in \mathbb{X}$, for $\varepsilon \in (0, \varepsilon'_0], i \in \mathbb{X}$, where $\varepsilon'_0 = \min_{j \in \mathbb{Y}_i} \varepsilon_{ij} =$

$\min_{j \in \mathbb{Y}_i, j \neq j_i} \varepsilon_{ij}$. Thus, condition **F** holds, if parameter ε_0 is replaced by the new value ε'_0 . In this case, functions $p_{ij}(\varepsilon), i, j \in \mathbb{X}$ can, for every $\varepsilon \in (0, \varepsilon'_0]$, serve as transition probabilities of a Markov chain.

Note that remainders $o_{ij}(\varepsilon^{l+}_{ij}), j \in \mathbb{Y}_i, i \in \mathbb{X}$ constructed above can be very irregular functions. Let us, for example, consider the case, where all asymptotic expansions in condition **D** have the same order, i.e., parameters $l^+_{ij} = l^+, j \in \mathbb{Y}_i, i \in \mathbb{X}$. In this case, identities (20) take the form, $\sum_{j \in \mathbb{Y}_i} o_{ij}(\varepsilon^{l+}) = 0, \varepsilon \in (0, \varepsilon_0], i \in \mathbb{X}$. Condition **D** requires that $o_{ij}(\varepsilon^{l+})/\varepsilon^{l+} \rightarrow 0$ as $\varepsilon \rightarrow 0$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$. Remainders $o_{ij}(\varepsilon^{l+}), j \in \mathbb{Y}_i, i \in \mathbb{X}$ can be continuous functions of ε taking zero value in at most finite numbers of points. However, let us multiply them, for example, by the Dirichlet function $D(\varepsilon)$. The new remainders $o'_{ij}(\varepsilon^{l+}) = D(\varepsilon)o_{ij}(\varepsilon^{l+}), j \in \mathbb{Y}_i, i \in \mathbb{X}$ also satisfy identities (20) and $o'_{ij}(\varepsilon^{l+})/\varepsilon^{l+} \rightarrow 0$ as $\varepsilon \rightarrow 0$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$. At the same time, they are very irregular functions. This example is, of course, an artificial one. But, it well illustrates the above statement about possible irregularity of remainders and, in sequel, transition probabilities, as functions of the perturbation parameter.

Let us also make some remarks concerned the expected sojourn times.

First, let us define $e_{ij}(\varepsilon) = 0, \varepsilon \in (0, \varepsilon_0], j \in \bar{\mathbb{Y}}_i, i \in \mathbb{X}$ that is consistent with condition **A (b)**.

Let us also $e_{ij}(\varepsilon), \varepsilon \in (0, \varepsilon_0], j \in \mathbb{Y}_i, i \in \mathbb{X}$ be some real-valued functions which satisfy condition **E**, i.e., can be represented in the form of Laurent asymptotic expansions, $e_{ij}(\varepsilon) = \sum_{l=m^-_{ij}}^{m^+_{ij}} b_{ij}[l]\varepsilon^l + \dot{o}_{ij}(\varepsilon^{m^+_{ij}}), \varepsilon \in (0, \varepsilon_0]$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$, where **(a)** $b_{ij}[m^-_{ij}] > 0$ and $-\infty < m^-_{ij} \leq m^+_{ij} < \infty$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$; **(b)** $\dot{o}_{ij}(\varepsilon^{m^+_{ij}})/\varepsilon^{m^+_{ij}} \rightarrow 0$ as $\varepsilon \rightarrow 0$, for $j \in \mathbb{Y}_i, i \in \mathbb{X}$.

Condition **E (b)** guarantees that, for every $j \in \mathbb{Y}_i, i \in \mathbb{X}$ there exists $\dot{\varepsilon}_{\alpha_{ij}, ij} \in (0, \varepsilon_0]$ such that, for $\varepsilon \in (0, \dot{\varepsilon}_{\alpha_{ij}, ij}]$,

$$|\dot{o}_{ij}(\varepsilon^{m^+_{ij}})/\varepsilon^{m^+_{ij}}| \leq \alpha_{ij}|b_{ij}[m^-_{ij}]|. \quad (24)$$

It is useful to note that in the case, where condition **E'** is assumed to hold, an explicit value for parameters $\dot{\varepsilon}_{\alpha_{ij}, ij}$ can be derived from the inequalities, $|\dot{o}_{ij}(\varepsilon^{m^+_{ij}})| \leq \dot{G}_{ij}\varepsilon^{m^+_{ij}+\delta_{ij}}, \varepsilon \in (0, \varepsilon_{ij}], j \in \mathbb{Y}_i, i \in \mathbb{X}$, penetrating this condition. These inequalities yield that relation (24) holds for $\dot{\varepsilon}_{\alpha_{ij}, ij} = \dot{\varepsilon}_{ij} \wedge \left(\frac{\alpha_{ij}|b_{ij}[m^-_{ij}]|}{\dot{G}_{ij}}\right)^{\frac{1}{\delta_{ij}}}, j \in \mathbb{Y}_i, i \in \mathbb{X}$.

Let us also define $B_{\varepsilon_0, ij} = \sum_{m_{ij}^- < l \leq m_{ij}^+} |b_{ij}[l]| \varepsilon_0^{l-m_{ij}^- - 1}$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$ and $\dot{\varepsilon}_{\alpha_{ij}, ij}'' = \dot{\varepsilon}_{\alpha_{ij}, ij}$ if $B_{\varepsilon_0, ij} = 0$ or $\dot{\varepsilon}_{\alpha_{ij}, ij}'' = \dot{\varepsilon}_{\alpha_{ij}, ij} \wedge \frac{\alpha_{ij} |b_{ij}[l_{ij}^-]|}{B_{\varepsilon_0, ij}}$ if $B_{\varepsilon_0, ij} > 0$.

The following inequality holds, for every $\varepsilon \in (0, \dot{\varepsilon}_{\alpha_{ij}, ij}'']$ and $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$,

$$\begin{aligned} e_{ij}(\varepsilon) &\geq \varepsilon^{m_{ij}^-} (b_{ij}[m_{ij}^-] - \varepsilon B_{\varepsilon_0, ij} - \varepsilon_0^{m_{ij}^+ - m_{ij}^-} |o_{ij}(\varepsilon^{m_{ij}^+}) / \varepsilon^{m_{ij}^+}|) \\ &\geq \varepsilon^{m_{ij}^-} b_{ij}[m_{ij}^-] (1 - 2\alpha_{ij}) > 0. \end{aligned} \quad (25)$$

Let us now define $\varepsilon_0'' = \min_{j \in \mathbb{Y}_i, i \in \mathbb{X}} \dot{\varepsilon}_{\alpha_{ij}, ij}''$.

Obviously, $e_{ij}(\varepsilon) > 0$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$, $\varepsilon \in (0, \varepsilon_0'']$. This is consistent, with conditions **A (a)** and **B**.

Finally, let us define $\tilde{\varepsilon}_0 = \varepsilon_0' \wedge \varepsilon_0''$. Parameter, $\tilde{\varepsilon}_0$ can serve as a new value for parameter ε_0 .

Functions $p_{ij}(\varepsilon)$, $i, j \in \mathbb{X}$ and $e_{ij}(\varepsilon)$, $i, j \in \mathbb{X}$ constructed above can serve, respectively, as transition probabilities of the embedded Markov chain $\eta_n^{(\varepsilon)}$ and expectations of sojourn times for some semi-Markov process $\eta^{(\varepsilon)}(t)$, for every $\varepsilon \in (0, \tilde{\varepsilon}_0]$. A variant of transition probabilities for such semi-Markov processes is given in Section 3*.

In conclusion, let us consider a numerical example.

We assume that the semi-Markov process $\eta^{(\varepsilon)}(t)$ is, for every $\varepsilon \in (0, \varepsilon_0]$, a semi-Markov process with the phase space $\mathbb{X} = \{1, 2, 3\}$.

The transition sets are $\mathbb{Y}_1 = \{1, 2\}$, $\mathbb{Y}_2 = \{1, 2, 3\}$, $\mathbb{Y}_3 = \{1, 2\}$.

The 3×3 matrix of transition probabilities $\|p_{ij}(\varepsilon)\|$, for the corresponding embedded Markov chain $\eta_n^{(\varepsilon)}$, has the following form,

$$\left\| \begin{array}{ccc} 1 - \varepsilon^2 - \varepsilon^3 + o_{11}(\varepsilon^3) & \varepsilon^2 + \varepsilon^3 + o_{12}(\varepsilon^3) & 0 \\ \frac{\varepsilon}{2} + \frac{\varepsilon^2}{2} - \varepsilon^3 + o_{21}(\varepsilon^3) & 1 - \varepsilon - \varepsilon^2 + o_{22}(\varepsilon^2) & \frac{\varepsilon}{2} + \frac{\varepsilon^2}{2} + 2\varepsilon^3 + o_{23}(\varepsilon^3) \\ \frac{1}{2} + \varepsilon^2 - \varepsilon^3 + o_{31}(\varepsilon^3) & \frac{1}{2} - \varepsilon^2 + \varepsilon^3 + o_{32}(\varepsilon^3) & 0 \end{array} \right\|. \quad (26)$$

The 3×3 matrix of expectations of sojourn times $\|e_{ij}(\varepsilon)\|$, for the semi-Markov process $\eta^{(\varepsilon)}(t)$, has the following form,

$$\left\| \begin{array}{ccc} \varepsilon + \varepsilon^2 + \dot{o}_{11}(\varepsilon^2) & \varepsilon^3 + \varepsilon^4 + \dot{o}_{12}(\varepsilon^4) & 0 \\ \varepsilon + \varepsilon^2 + \dot{o}_{21}(\varepsilon^2) & 1 - \varepsilon + \varepsilon^3 + \dot{o}_{22}(\varepsilon^3) & 2 + \varepsilon + \varepsilon^2 + \dot{o}_{23}(\varepsilon^2) \\ \varepsilon^{-1} + 1 + \dot{o}_{31}(1) & 2\varepsilon^{-1} + \varepsilon + \dot{o}_{32}(\varepsilon) & 0 \end{array} \right\|. \quad (27)$$

In the asymptotic expansions penetrating relations (26) and (27), the coefficients $a_{ij}[l_{ij}^-]$, $b_{ij}[m_{ij}^-] > 0$, $j \in \mathbb{Y}_i$, $i \in \mathbb{X}$, and coefficients $a_{ij}[l]$, $l =$

$l_{ij}^-, \dots, l_{ij}^+, j \in \mathbb{Y}_i, i \in \mathbb{X}$ satisfy relation (19). We also assume that parameter $\varepsilon_0 = \tilde{\varepsilon}_0$ and remainders $o(\varepsilon_{ij}^{l_{ij}^+}), \dot{o}(\varepsilon_{ij}^{m_{ij}^+}), j \in \mathbb{Y}_i, i \in \mathbb{X}$, in the asymptotic expansions representing elements of matrices given in relations (26) and (27), are chosen according the procedures described above, in particular, the identities (20) hold. In this case, matrices, given in the above relations, can, for every $\varepsilon \in (0, \varepsilon_0]$, serve as, respectively, the matrix of transition probabilities for the corresponding embedded Markov chain and the matrix of expectations of sojourn times, for the semi-Markov process $\eta^{(\varepsilon)}(t)$, and conditions **A** – **E** hold.

The matrices of transition probabilities $\|p_{ij}(\varepsilon)\|$, for the embedded Markov chains $\eta_n^{(\varepsilon)}$, and $\|p_{ij}(0)\|$, for the limiting Markov chain $\eta_n^{(0)}$, have, respectively, the following forms,

$$\left\| \begin{array}{ccc} \bullet & \bullet & 0 \\ \bullet & \bullet & \bullet \\ \bullet & \bullet & 0 \end{array} \right\| \quad \text{and} \quad \left\| \begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 \end{array} \right\|, \quad (28)$$

where symbol $\bullet$ indicates positions of positive elements in matrices $\|p_{ij}(\varepsilon)\|$, $\varepsilon \in (0, \varepsilon_0]$.

The phase space $\mathbb{X}$ is one class of communicative states for the embedded Markov chain $\eta_n^{(\varepsilon)}$, for every $\varepsilon \in (0, \varepsilon_0]$, while it consists of two closed classes of communicative states $\mathbb{X}_1 = \{1\}$, $\mathbb{X}_2 = \{2\}$ and the class of transient states $\mathbb{X}_3 = \{3\}$, for the limiting Markov chain $\eta_n^{(0)}$.

By excluding the state 1 from the phase space $\mathbb{X}$ and using the algorithm described in Section 5*, we construct the reduced semi-Markov processes ${}_1\eta^{(\varepsilon)}(t)$, with the phase space ${}_1\mathbb{X} = \{2, 3\}$. Conditions **A** – **E** hold for these reduced semi-Markov processes. The corresponding transition sets are ${}_1\mathbb{Y}_2 = \{2, 3\}$ and ${}_1\mathbb{Y}_3 = \{2\}$. By applying the algorithms described in Lemma 8* and Theorems 2* and 3*, we can compute the 2×2 matrices ${}_1\mathbf{P} = \|{}_1p_{ij}(\varepsilon)\|$ and ${}_1\mathbf{E} = \|{}_1e_{ij}(\varepsilon)\|$. These matrices take the following forms,

$${}_1\mathbf{P} = \left\| \begin{array}{cc} 1 - \frac{1}{2}\varepsilon - \frac{1}{2}\varepsilon^2 + {}_1o_{22}(\varepsilon^2) & \frac{\varepsilon}{2} + \frac{\varepsilon^2}{2} + 2\varepsilon^3 + {}_1o_{23}(\varepsilon^3) \\ 1 & 0 \end{array} \right\|, \quad (29)$$

and

$${}_1\mathbf{E} = \left\| \begin{array}{cc} \frac{3}{2} + \frac{1}{2}\varepsilon + {}_1\dot{o}_{22}(\varepsilon) & 2 + \varepsilon + \varepsilon^2 + {}_1\dot{o}_{23}(\varepsilon^2) \\ \frac{7}{2}\varepsilon^{-1} + 1 + {}_1\dot{o}_{32}(1) & 0 \end{array} \right\|. \quad (30)$$

By excluding the state 2 from the reduced phase space ${}_1\mathbb{X} = \{2, 3\}$, we construct the “final” reduced semi-Markov processes ${}_{(1,2)}\eta^{(\varepsilon)}(t)$, with the

one-state phase space ${}_{(1,2)}\mathbb{X} = \{3\}$. Conditions **A** – **E** also hold for these semi-Markov processes. The corresponding transition set ${}_{(1,2)}\mathbb{Y} = \{3\}$. The transition probability ${}_{(1,2)}p_{33}(\varepsilon) \equiv 1$. By applying the algorithms described in Theorem 4_{*}, we can also compute the Laurent asymptotic expansion for the expected return time, ${}_{(1,2)}e_{33}(\varepsilon) = E_{33}(\varepsilon) = \frac{21}{2}\varepsilon^{-1} - 3 + \ddot{o}_{33}(1)$. The Laurent asymptotic expansion for the expected sojourn time $e_3(\varepsilon) = e_{31}(\varepsilon) + e_{32}(\varepsilon) + e_{33}(\varepsilon)$, obtained with the use of the multiple summation rule given in Lemma 3_{*}, has the following form, $e_3(\varepsilon) = 3\varepsilon^{-1} + 1 + \dot{o}_3(1)$. Finally, the algorithm described in Theorem 5_{*} gives the following asymptotic expansion, for the stationary probability $\pi_3(\varepsilon) = \frac{e_3(\varepsilon)}{E_{33}(\varepsilon)} = \frac{3\varepsilon^{-1} + 1 + \dot{o}_3(1)}{\frac{21}{2}\varepsilon^{-1} - 3 + \ddot{o}_{33}(1)} = \frac{2}{7} + \frac{26}{147}\varepsilon + o_3(\varepsilon)$.

Also, by excluding the state 3 from the reduced phase space ${}_1\mathbb{X} = \{2, 3\}$ and applying the algorithms described in Theorem 4_{*}, we can compute the Laurent asymptotic expansion for the expected return time ${}_{(1,3)}e_{22}(\varepsilon) = E_{22}(\varepsilon) = \frac{21}{4} + \frac{15}{4}\varepsilon + \ddot{o}_{22}(\varepsilon)$. In this case, the asymptotic expansion for the expected sojourn time $e_2(\varepsilon) = 3 + \varepsilon + 2\varepsilon^2 + \dot{o}_2(\varepsilon^2)$, and the algorithm described in Theorem 5_{*} gives the following asymptotic expansion for the stationary probability $\pi_2(\varepsilon) = \frac{e_2(\varepsilon)}{E_{22}(\varepsilon)} = \frac{3 + \varepsilon + 2\varepsilon^2 + \dot{o}_2(\varepsilon^2)}{\frac{21}{4} + \frac{15}{4}\varepsilon + \ddot{o}_{22}(\varepsilon)} = \frac{4}{7} - \frac{32}{147}\varepsilon + o_2(\varepsilon)$.

As far as the stationary probability $\pi_1(\varepsilon)$ is concerned, the corresponding asymptotic expansion can be found using the identity, $\pi_1(\varepsilon) = 1 - \pi_2(\varepsilon) - \pi_3(\varepsilon)$, $\varepsilon \in (0, \varepsilon_0]$, and the operational rules for asymptotic expansions given in Lemma 3_{*}. This yields the asymptotic expansion, $\pi_1(\varepsilon) = \frac{1}{7} + \frac{6}{147}\varepsilon + o_1(\varepsilon)$.

Alternatively, the exclusion of states from the phase space $\mathbb{X}$ in the opposite order, first state 3 and then state 2 or 1, and the use of the algorithms described in Lemma 8_{*} and Theorems 2_{*} – 4_{*} yield the asymptotic expansions for the expected return times ${}_{(3,2)}e_{11}(\varepsilon) = E_{11}(\varepsilon) = 7\varepsilon + \frac{15}{3}\varepsilon^2 + \ddot{o}_{11}(\varepsilon^2)$ and ${}_{(3,1)}e_{22}(\varepsilon) = E_{22}(\varepsilon) = \frac{21}{4} + \frac{15}{4}\varepsilon + \ddot{o}_{22}(\varepsilon)$. Then, the algorithm described in Theorem 5_{*} yields the same asymptotic expansions for stationary probabilities $\pi_1(\varepsilon) = \frac{e_1(\varepsilon)}{E_{11}(\varepsilon)} = \frac{\varepsilon + \varepsilon^2 + \dot{o}_1(\varepsilon^2)}{7\varepsilon + \frac{15}{3}\varepsilon^2 + \ddot{o}_{11}(\varepsilon^2)} = \frac{1}{7} + \frac{6}{147}\varepsilon + o_1(\varepsilon)$ and $\pi_2(\varepsilon) = \frac{e_2(\varepsilon)}{E_{22}(\varepsilon)} = \frac{3 + \varepsilon + 2\varepsilon^2 + \dot{o}_2(\varepsilon^2)}{\frac{21}{4} + \frac{15}{4}\varepsilon + \ddot{o}_{22}(\varepsilon)} = \frac{4}{7} - \frac{32}{147}\varepsilon + o_2(\varepsilon)$.

Note that the Laurent asymptotic expansion for the expectation $E_{22}(\varepsilon)$ and, in sequel, the Taylor asymptotic expansion for the stationary probability $\pi_2(\varepsilon)$ are invariant with respect to the choice the sequence of states $\langle 1, 3 \rangle$ or $\langle 3, 1 \rangle$ for sequential exclusion from the phase space $\mathbb{X}$. This is consistent with the corresponding invariance statements formulated in Theorems 4_{*} and 5_{*}.

The coefficients of the asymptotic expansions $\pi_i(\varepsilon) = c_i[0] + c_i[1]\varepsilon + o_i(\varepsilon)$, $i = 1, 2, 3$ given above satisfy relations, $c_1[0] + c_2[0] + c_3[0] = 1$ and

$c_1[1] + c_2[1] + c_3[1] = 0$. This is consistent with the corresponding statement in Theorem 5*.

In the example presented above, we did not trace the explicit formulas for remainders $o_1(\varepsilon)$, $o_2(\varepsilon)$ and $o_3(\varepsilon)$. However, according to the corresponding statement in Theorem 5*, these remainders are connected by the following identity $o_1(\varepsilon) + o_2(\varepsilon) + o_3(\varepsilon) \equiv 0$.

We would like also to explain an unexpected, in some sense, asymptotic behavior of stationary probabilities $\pi_i(\varepsilon)$, in the above example. As a matter of fact, states 1 and 2 are asymptotically absorbing states with non-absorption probabilities of different order, respectively, $O(\varepsilon^2)$ and $O(\varepsilon)$. While, state 3 is a transient asymptotically non-absorbing state. This, seems, should cause convergence of the stationary probability $\pi_1(\varepsilon)$ to 1 and the stationary probabilities $\pi_2(\varepsilon)$ and $\pi_3(\varepsilon)$ to 0 as $\varepsilon \rightarrow 0$, with different rates of convergence. This, however, does not take place, and all three probabilities converge to non-zero limits. This is because of the expected sojourn times $e_1(\varepsilon)$, $e_2(\varepsilon)$ and $e_3(\varepsilon)$ have orders, respectively, $O(\varepsilon)$, $O(1)$ and $O(\varepsilon^{-1})$. These expectations compensate absorption effects for states 1, 2 and 3.

In the above example, computations of explicit upper bounds for remainders in the asymptotic expansions for stationary probabilities $\pi_1(\varepsilon)$, $\pi_2(\varepsilon)$ and $\pi_3(\varepsilon)$ can also be realized in the case, where conditions **D'** and **E'** hold instead of conditions **D** and **E**. We, however, omit this presentation, in order to escape overloading the paper by technical numerical computations.

References

- [1] AVRACHENKOV, K. E., FILAR, J. A. AND HOWLETT, P. G. (2013). *Analytic Perturbation Theory and Its Applications*. SIAM, Philadelphia, PA, xii+372 pp.
- [2] BINI, D. A., LATOUCHE, G. AND MEINI, B. (2005). *Numerical Methods for Structured Markov Chains*. Numerical Mathematics and Scientific Computation, Oxford Science Publications, Oxford University Press, New York, xii+327 pp.
- [3] COURTOIS, P. J. (1977). *Decomposability. Queueing and Computer System Applications*. ACM Monograph Series, Academic Press, New York, xiii+201 pp.
- [4] GYLLENBERG, M. AND SILVESTROV, D. S. (2008). *Quasi-Stationary Phenomena in Nonlinearly Perturbed Stochastic Systems*. De Gruyter Expositions in Mathematics, **44**, Walter de Gruyter, Berlin, ix+579 pp.
- [5] KARTASHOV, M. V. (1996). *Strong Stable Markov Chains*. VSP, Utrecht and TBiMC, Kiev, 138 pp.

- [6] KONSTANTINOV, M., GU, D. W., MEHRMANN, V. AND PETKOV, P. (2003). *Perturbation Theory for Matrix Equations*. Studies in Computational Mathematics, **9**, North-Holland, Amsterdam, xii+429 pp.
- [7] KOROLYUK, V. S. AND KOROLYUK, V. V. (1999). *Stochastic Models of Systems*. Mathematics and its Applications, **469**, Kluwer, Dordrecht, xii+185 pp.
- [8] KOROLIUK, V. S. AND LIMNIOS, N. (2005). *Stochastic Systems in Merging Phase Space*. World Scientific, Singapore, xv+331 pp.
- [9] KOROLYUK, V. S. AND TURBIN, A. F. (1976). *Semi-Markov Processes and its Applications*. Naukova Dumka, Kiev, 184 pp.
- [10] KOROLYUK, V. S. AND TURBIN, A. F. (1978). *Mathematical Foundations of the State Lumping of Large Systems*. Naukova Dumka, Kiev, 218 pp. (English edition: Mathematics and its Applications, **264**, Kluwer, Dordrecht, 1993, x+278 pp.).
- [11] SENETA, E. (2006). *Non-Negative Matrices and Markov Chains*. Springer Series in Statistics. Springer, New York, xvi+287 pp. (A revised reprint of the second (1981) edition).
- [12] SILVESTROV, D. AND SILVESTROV, S. (2015). Asymptotic expansions for stationary distributions of perturbed semi-Markov processes. Research Report 2015-9, Department of Mathematics, Stockholm University, 75 pp. and arXiv:1603.03891.
- [13] SILVESTROV, D. AND SILVESTROV, S. (2016). Asymptotic expansions for stationary distributions of nonlinearly perturbed semi-Markov processes. I.
- [14] STEWART, G. W. (1998). *Matrix Algorithms. Vol. I. Basic Decompositions*. SIAM, Philadelphia, PA, xx+458 pp.
- [15] STEWART, G. W. (2001). *Matrix Algorithms. Vol. II. Eigensystems*. SIAM, Philadelphia, PA, xx+469 pp.
- [16] STEWART, G. W. AND SUN, J. G. (1990). *Matrix Perturbation Theory*. Computer Science and Scientific Computing. Academic Press, Boston, xvi+365 pp.
- [17] YIN, G. G. AND ZHANG, Q. (2005). *Discrete-Time Markov Chains. Two-Time-Scale Methods and Applications*. Stochastic Modelling and Applied Probability, **55**, Springer, New York, xix+348 pp.
- [18] YIN, G. G. AND ZHANG, Q. (2013). *Continuous-Time Markov Chains and Applications. A Two-Time-Scale Approach*. Second edition, Stochastic Modelling and Applied Probability, **37**, Springer, New York, xxii+427 pp. (An extended variant of the first (1998) edition).